(19) Europäisches Patentamt
European Patent Office
Office européen des brevets

(11) Publication number: 0 346 710 A2

(12) EUROPEAN PATENT APPLICATION

(21) Application number: 89110096.8

(22) Date of filing: 03.06.89

(51) Int. Cl.4: C12N 15/00 , C12N 5/00 , C07K 13/00 , A61K 39/395

The microorganism(s) has (have) been deposited with ATCC under number(s) Nos 67708,67709,67710,67711,67712.

(30) Priority: 16.06.88 US 207678
21.11.88 US 274107

(43) Date of publication of application:
20.12.89 Bulletin 89/51

(84) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI NL SE

(71) Applicant: MOLECULAR DIAGNOSTICS, INC.
400 Morgan Lane
West Haven, CT 06516(US)

(72) Inventor: Barnett, Thomas R., Dr.
27 Jeffrey Road
East Haven, CT 06513(US)
Inventor: Elting, James J., Dr.
5 Heatherwood Drive
Madison, CT 06443(US)
Inventor: Kamarck, Michael E.
86 Russell Road
Bethany, CT 06525(US)
Inventor: Kretschmer, Axel, Dr.
Richard-Zörner-Strasse 32
D-5060 Bergisch Gladbach 1(DE)

(74) Representative: Adrian, Albert, Dr. et al
c/o BAYER AG Konzernverwaltung RP
Patentabteilung
D-5090 Leverkusen 1, Bayerwerk(DE)

(54) cDNAs coding for members of the carcinoembryonic antigen family.

(57) A nucleic acid comprising a base sequence which codes for a CEA family member peptide sequence or nucleic acids having a base sequence hybridizable therewith, replicable recombinant cloning vehicles having an insert comprising such nucleic acid, cells transfected, infected or injected with such cloning vehicles, polypeptides expressed by such cells, synthetic peptides derived from the coding sequence of CEA family member nucleic acids, antibody preparations specific for such polypeptides, immunoassays for detecting CEA family members using such antibody preparations and nucleic acid hybridization methods for detecting CEA family member nucleic acid sequences using a nucleic acid probe comprising the above described nucleic acid.

EP 0 346 710 A2

cDNAs coding for members of the carcinoembryonic antigen family

BACKGROUND OF THE INVENTION

Field of the Invention

The present invention concerns nucleic acid sequences which code for carcinoembryonic antigen (CEA) antigen family peptide sequences.

Background Information

Carcinoembryonic antigen was first described by Gold and Freedman, J. Exp. Med., 121, 439-462, (1965). CEA is characterized as a glycoprotein of approximately 200,000 molecular weight with 50-60% by weight of carbohydrate. CEA is present during normal human fetal development, but only in very low concentration in the normal adult intestinal tract. It is produced and secreted by a number of different tumors.

CEA is a clinically useful tumor marker for the management of colorectal cancer patients. CEA can be measured using sensitive immunoassay methods. When presurgical serum levels of CEA are elevated, a postsurgical drop in serum CEA to the normal range typically indicates successful resection of the tumor. Postsurgical CEA levels that do not return to normal often indicate incomplete resection of the tumor or the presence of additional tumor sites in the patient. After returning to normal levels, subsequent rapid rises in serum CEA levels usually indicate the presence of metastages. Slower postsurgical rises from the normal level are most often interpreted to indicate the presence of new primary tumors not previously detected. Post surgical management of colon cancer patients is thus facilitated by the measurement of CEA.

CEA is a member of an antigen family. Because of this, the immunoassay of CEA by presently available methods is complicated by the fact that CEA is but one of several potentially reactive antigens. There have been at least sixteen CEA-like antigens described in the literature. Since some of these appear to be the same antigen described by different investigators, the actual number of different antigens is somewhat less than this number. Nonetheless, there is a complex array of cross-reactive antigens which can potentially interfere with an immunoassay of the CEA released by tumors. It is known that serum levels of CEA-like antigens are elevated in many non-cancerous conditions such an inflammatory liver diseases and also in smokers. It is important that immunoassays used for the monitoring of cancer patient status not be interfered with by these other CEA-like antigens. Conversely, it is important to be able to distinguish the antigens by immunoassays because of the possibility that different tumor types may preferentially express different forms of CEA. If so, then the ability to reliably measure the different forms of CEA can provide the means to diagnose or more successfully treat different forms of cancer.

The members of the "CEA family" share some antigenic determinants. These common epitopes are not useful in distinguishing the members of the antigen family and antibodies recognizing them are of little use for measuring tumor-specific CEA levels.

U.S.P. 3,663,684, entitled "Carcinoembryonic Antigen and Diagnostic Method using Radioactive Iodine", concerns purification and radioiodination of CEA for use in a RIA.

U.S.P. 3,697,638 describes that CEA is a mixture of antigens (components A and B in this case). U.S.P. 3,697,638 mentions methods for separating and radioiodinating each component and their use in specific RIA's.

U.S.P. 3,852,415, entitled "Compositions for Use in Radioimmunoassay, as Substitute for Blood Plasma Extract in Determination of Carcinoembryonic Antigen" relates to the use of a buffer containing EDTA and bovine serum albumin as a substitute for plasma as a diluent for CEA RIA's.

U.S.P. 3,867,363, entitled "Carcinoembryonic Antigens", is directed to the isolation of CEA components A and B, their labelling and use in a RIA.

U.S.P. 3,927,193, entitled "Localization of Tumors by Radiolabelled Antibodies", concerns the use of radiolabelled anti-CEA antibodies in whole body tumor imaging.

U.S.P. 3,956,258, entitled "Carcinoembryonic Antigens", relates to the isolation of CEA components A and B.

U.S.P. 4,086,217, entitled "Carcinoembryonic Antigens", is directed to the isolation of CEA components

A and B.

U.S.P. 4,140,753, entitled "Diagnostic Method and Reagent", concerns the purification of a CEA isomer called CEA-S1 and its use in a RIA.

U.S.P. 4,145,336, entitled "Carcinoembryonic Antigen Isomer", relates to the antigen CEA-S1.

U.S.P. 4,180,499, entitled "Carcinoembryonic Antigens", describes a process for producing CEA component B.

U.S.P. 4,228,236, entitled "Process of Producing Carcinoembryonic Antigen", is directed to the use of the established cell lines LS-174T and LS-180 or clones or derivatives thereof for the production of CEA.

U.S.P. 4,272,504, entitled "Antibody Absorbed Support Method for Carcinoembryonic Antigen Assay", concerns two concepts for the radioimmunoassay of CEA. First, U.S.P. 4,272,504 relates to a sample pretreatment in the form of heating to 65 to 85° C at pH 5 to precipitate and eliminate extraneous protein. Second, it describes the use of a solid phase antibody (either on beads or tubes) as a means to capture analyte and radiolabelled CEA tracer.

U.S.P. 4,299,815, entitled "Carcinoembryonic Antigen Determination", concerns diluting a CEA sample with water and pretreating by heating to a temperature below which precipitation of protein will occur. The pretreated sample is then immunoassayed using RIA, EIA, FIA or chemiluminescent immunoassay.

U.S.P. 4,349,528, entitled "Monoclonal Hybridoma Antibody Specific for High Molecular Weight Carcinoembryonic Antigen", is directed to a monoclonal antibody reacting with 180 kD CEA, but not with other molecular weight forms.

U.S.P. 4,467,031, entitled "Enzyme-Immunoassay for Carcinoembryonic Antigen", relates to a sandwich enzyme immunoassay for CEA in which the first of two anti-CEA monoclonal antibodies is attached to a solid phase and the second monoclonal is conjugated with peroxidase.

U.S.P. 4,489,167, entitled "Methods and Compositions for Cancer Detection", describes that CEA shares an antigenic determinant with alpha-acid glycoprotein (AG), which is a normal component of human serum. The method described therein concerns a solid-phase sandwich enzyme immunoassay using as one antibody an antibody recognizing AG and another antibody recognizing CEA, but not AG.

U.S.P. 4,578,349, entitled "Immunoassay for Carcinoembryonic Antigen (CEA)", is directed to the use of high salt containing buffers as diluents in CEA immunoassays.

EP 113072-A, entitled "Assaying Blood Sample for Carcinoembryonic Antigen - After Removal of Interfering Materials by Incubation with Silica Gel", relates to the removal from a serum of a plasma sample of interfering substances by pretreatment with silica gel. The precleared sample is then subjected to an immunoassay.

EP 102008-A, entitled "Cancer Diagnostics Carcinoembryonic Antigen - Produced from Perchloric Acid Extracts Without Electrophoresis", relates to a procedure for the preparation of CEA from perchloric acid extracts, without the use of an electrophoresis step.

EP 92223-A, entitled "Determination of Carcinoembryonic Antigen in Cytosol or Tissue - for Therapy Control and Early Recognition of Regression", concerns an immunoassay of CEA, not in serum or plasma, but in the cytosol fraction of the tumor tissue itself.

EP 83103759.6, entitled "Cytosole-CEA-Measurement as Predictive Test in Carcinoma, Particularly Mammacarcinoma", is similar to EP 92223-A.

EP 83303759, entitled "Monoclonal Antibodies Specific to Carcinoembryonic Antigen", relates to the production of "CEA specific" monoclonal antibodies and their use in immunoassays.

WO 84/02983, entitled "Specific CEA-Family Antigens, Antibodies Specific Thereto and Their Methods of Use", is directed to the use of monoclonal antibodies to CEA-meconium (MA)-, and NCA-specific epitopes in immunoassays designed to selectively measure each of these individual components in a sample.

All of the heretofore CEA assays utilize either monoclonal or polyclonal antibodies which are generated by immunizing animals with the intact antigen of choice. None of them address the idea of making sequence specific antibodies for the detection of a unique primary sequence of the various antigens. They do not cover the use of any primary amino acid sequence for the production of antibodies to synthetic peptides or fragments of the natural product. They do not include the concept of using primary amino acid sequences to distinguish the CEA family members. None of them covers the use of DNA or RNA clones for isolating the genes with which to determine the primary sequence.

DEFINITIONS

Nucleic Acid Abbreviations	
A	adenine
G	guanine
C	cytosine
T	thymidine
U	uracil

Amino Acid Abbreviations:	
Asp	aspartic acid
Asn	asparagine
Thr	threonine
Ser	serine
Glu	glutamic acid
Gln	glutamine
Pro	proline
Gly	glycine
Ala	alanine
Cys	cysteine
Val	valine
Met	methionine
Ile	isoleucine
Leu	leucine
Tyr	tyrosine
Phe	phenylalanine
Trp	tryptophan
Lys	lysine
His	histidine
Arg	arginine

Nucleotide - A monomeric unit of DNA or RNA containing a sugar moiety (pentose), a phosphate, and a nitrogenous heterocyclic base. The base is linked to the sugar moiety via the glycosidic carbon (1′ carbon of the pentose) and that combination of base and sugar is called a nucleoside. The base characterizes the nucleotide. The four DNA bases are adenine ("A"), guanine ("G"), cytosine ("C"), and thymine ("T"). The four RNA bases are A, G, C and uracil ("U").

DNA Sequence - A linear array of nucleotides connected one to the other by phosphodiester bonds between the 3′ and 5′ carbons of adjacent pentoses.

Functional equivalents - It is well known in the art that in a DNA sequence some nucleotides can be replaced without having an influence on the sequence of the expression product. With respect to the peptide this term means that one or more amino acids which have no function in a particular use can be deleted or replaced by another one.

Codon - A DNA sequence of three nucleotides (a triplet) which encodes through mRNA an amino acid, a translation start signal or a translation termination signal. For example, the nucleotide triplets TTA, TTG, CTT, CTC, CTA and CTG encode the amino acid leucine ("Leu"), TAG, TAA and TGA are translation stop signals and ATG is a translation start signal.

Reading Frame - The grouping of codons during translation of mRNA into amino acid sequences. During translation, the proper reading frame must be maintained. For example, the sequence GCTGGTTGTAAG may be translated in three reading frames or phases, each of which affords a different amino acid sequence

```
GCT GGT TGT AAG   - Ala-Gly-Cys-Lys
G CTG GTT GTA AG  - Leu-Val-Val
GC TGG TTG TAA G  - Trp-Leu-(STOP).
```

Polypeptide - A linear array of amino acids connected one to the other by peptide bonds between the alpha-amino and carboxy groups of adjacent amino acids.

Genome - The entire DNA of a cell or a virus. It includes inter alia the structural genes coding for the polypeptides of the cell or virus, as well as its operator, promoter and ribosome binding and interaction sequences, including sequences such as the Shine-Dalgarno sequences.

Structural Gene - A DNA sequence which encodes through its template or messenger RNA ("mRNA") a sequence of amino acids characteristic of a specific polypeptide.

Transcription - The process of producing mRNA from a structural gene.

Translation - The process of producing a polypeptide from mRNA.

Expression - The process undergone by a structural gene to produce a polypeptide. It is a combination of transcription and translation.

Plasmid - A non-chromosomal double-stranded DNA sequence comprising an intact "replicon" such that the plasmid is replicated in a host cell. When the plasmid is placed within a unicellular organism, the characteristics of that organism may be changed or transformed as a result of the DNA of the plasmid. For example, a plasmid carrying the gene for tetracycline resistance (Tet^R) transforms a cell previously sensitive to tetracycline into one which is resistant to it. A cell transformed by a plasmid is called a "transformant".

Phage or Bacteriophage - Bacterial virus, many of which consist of DNA sequences encapsulated in a protein envelope or coat ("capsid protein").

Cloning Vehicle - A plasmid, phage DNA or other DNA sequence which is capable of replicating in a host cell, which is characterized by one or a small number of endonuclease recognition sites at which such DNA sequences may be cut in a determinable fashion without attendant loss of an essential biological function of the DNA, e.g., replication, production of coat proteins or loss of promoter or binding sites, and which contains a marker suitable for use in the identification of transformed cells, e.g., tetracycline resistance or ampicillin resistance. A cloning vehicle is often called a vector.

Cloning - The process of obtaining a population of organisms or DNA sequences derived from one such organism or sequence by asexual reproduction.

Recombinant DNA Molecule or Hybrid DNA - A molecule consisting of segments of DNA from different genomes which have been joined end-to-end outside of living cells and have the capacity to infect some host cell and be maintained therein.

cDNA Expression Vector - A procaroytic cloning vehicle which also contains sequences of nucleotides that facilitate expression of cDNA sequences in eucaroytic cells. These nucleotides include sequences that function as eucaryotic promoter, alternative splice sites and polyadenylation signals.

Transformation/Transfection - DNA or RNA is introduced into cells in such a way as to allow gene expression. "Infected" referred to herein concerns the introduction of RNA or DNA by a viral vector into the host.

"Injected" referred to herein concerns the microinjection (use of a small syringe) of DNA into a cell.

CEA antigen family (CEA gene family) - a set of genes (gene family) and their products (antigen family) that share nucleotide sequences homologous to partial cDNA LV-7 (CEA-(a)) and as a result of theses similarities also share a subset of their antigenic epitopes. Examples of the CEA antigen family include CEA (= CEA-(b)), transmembrane CEA (TMCEA) = CEA-(c) and normal crossreacting antigen NCA (= CEA-(d)).

SUMMARY OF THE INVENTION

The present invention concerns the following DNA sequences designated as TM-2 (CEA-(e)), TM-3 (CEA-(f)), TM-4 (CEA-(g)), KGCEA1 and KGCEA2, which code for CEA antigen family peptide sequences or nucleic acids having a base sequence (DNA or RNA) that are hybridizable therewith:

SEQUENCE AND TRANSLATION OF cDNA OF TM-2

```
          10                  30                  50
CAGCCGTGCTCGAAGCGTTCCTGGAGCCCAAGCTCTCCTCCACAGGTGAAGACAGGGCCA

          70                  90                 110
GCAGGAGACACCATGGGGCACCTCTCAGCCCCACTTCACAGAGTGCGTGTACCCTGGCAG
              MetGlyHisLeuSerAlaProLeuHisArgValArgValProTrpGln

         130                 150                 170
GGGCTTCTGCTCACAGCCTCACTTCTAACCTTCTGGAACCCGCCCACCACTGCCCAGCTC
GlyLeuLeuLeuThrAlaSerLeuLeuThrPheTrpAsnProProThrThrAlaGlnLeu

         190                 210                 230
ACTACTGAATCCATGCCATTCAATGTTGCAGAGGGGAAGGAGGTTCTTCTCCTTGTCCAC
ThrThrGluSerMetProPheAsnValAlaGluGlyLysGluValLeuLeuLeuValHis

         250                 270                 290
AATCTGCCCCAGCAACTTTTTGGCTACAGCTGGTACAAAGGGGAAAGAGTGGATGGCAAC
AsnLeuProGlnGlnLeuPheGlyTyrSerTrpTyrLysGlyGluArgValAspGlyAsn

         310                 330                 350
CGTCAAATTGTAGGATATGCAATAGGAACTCAACAAGCTACCCCAGGGCCCGCAAACAGC
ArgGlnIleValGlyTyrAlaIleGlyThrGlnGlnAlaThrProGlyProAlaAsnSer

         370                 390                 410
GGTCGAGAGACAATATACCCCAATGCATCCCTGCTGATCCAGAACGTCACCCAGAATGAC
GlyArgGluThrIleTyrProAsnAlaSerLeuLeuIleGlnAsnValThrGlnAsnAsp

         430                 450                 470
ACAGGATTCTACACCCTACAAGTCATAAAGTCAGATCTTGTGAATGAAGAAGCAACTGGA
ThrGlyPheTyrThrLeuGlnValIleLysSerAspLeuValAsnGluGluAlaThrGly
```

```
          490                 510                 530
CAGTTCCATGTATACCCGGAGCTGCCCAAGCCCTCCATCTCCAGCAACAACTCCAACCCT
GlnPheHisValTyrProGluLeuProLysProSerIleSerSerAsnAsnSerAsnPro

          550                 570                 590
GTGGAGGACAAGGATGCTGTGGCCTTCACCTGTGAACCTGAGACTCAGGACACAACCTAC
ValGluAspLysAspAlaValAlaPheThrCysGluProGluThrGlnAspThrThrTyr

          610                 630                 650
CTGTGGTGGATAAACAATCAGAGCCTCCCGGTCAGTCCCAGGCTGCAGCTGTCCAATGGC
LeuTrpTrpIleAsnAsnGlnSerLeuProValSerProArgLeuGlnLeuSerAsnGly

          670                 690                 710
AACAGGACCCTCACTCTACTCAGTGTCACAAGGAATGACACAGGACCCTATGAGTGTGAA
AsnArgThrLeuThrLeuLeuSerValThrArgAsnAspThrGlyProTyrGluCysGlu

          730                 750                 770
ATACAGAACCCAGTGAGTGCGAACCGCAGTGACCCAGTCACCTTGAATGTCACCTATGGC
IleGlnAsnProValSerAlaAsnArgSerAspProValThrLeuAsnValThrTyrGly

          790                 810                 830
CCGGACACCCCCACCATTTCCCCTTCAGACACCTATTACCGTCCAGGGGCAAACCTCAGC
ProAspThrProThrIleSerProSerAspThrTyrTyrArgProGlyAlaAsnLeuSer

          850                 870                 890
CTCTCCTGCTATGCAGCCTCTAACCCACCTGCACAGTACTCCTGGCTTATCAATGGAACA
LeuSerCysTyrAlaAlaSerAsnProProAlaGlnTyrSerTrpLeuIleAsnGlyThr

          910                 930                 950
TTCCAGCAAAGCACACAAGAGCTCTTTATCCCTAACATCACTGTGAATAATAGTGGATCC
PheGlnGlnSerThrGlnGluLeuPheIleProAsnIleThrValAsnAsnSerGlySer

          970                 990                 1010
TATACCTGCCACGCCAATAACTCAGTCACTGGCTGCAACAGGACCACAGTCAAGACGATC
TyrThrCysHisAlaAsnAsnSerValThrGlyCysAsnArgThrThrValLysThrIle
```

1030 1050 1070

ATAGTCACTGATAATGCTCTACCACAAGAAAATGGCCTCTCACCTGGGGCCATTGCTGGC
IleValThrAspAsnAlaLeuProGlnGluAsnGlyLeuSerProGlyAlaIleAlaGly

1090 1110 1130

ATTGTGATTGGAGTAGTGGCCCTGGTTGCTCTGATAGCAGTAGCCCTGGCATGTTTTCTG
IleValIleGlyValValAlaLeuValAlaLeuIleAlaValAlaLeuAlaCysPheLeu

1150 1170 1190

CATTTCGGGAAGACCGGCAGGGCAAGCGACCAGCGTGATCTCACAGAGCACAAACCCTCA
HisPheGlyLysThrGlyArgAlaSerAspGlnArgAspLeuThrGluHisLysProSer

1210 1230 1250

GTCTCCAACCACACTCAGGACCACTCCAATGACCCACCTAACAAGATGAATGAAGTTACT
ValSerAsnHisThrGlnAspHisSerAsnAspProProAsnLysMetAsnGluValThr

1270 1290 1310

TATTCTACCCTGAACTTTGAAGCCCAGCAACCCACACAACCAACTTCAGCCTCCCCATCC
TyrSerThrLeuAsnPheGluAlaGlnGlnProThrGlnProThrSerAlaSerProSer

1330 1350 1370

CTAACAGCCACAGAAATAATTTATTCAGAAGTAAAAAAAGCAGTAATGAAACCTGTCCTGC
LeuThrAlaThrGluIleIleTyrSerGluValLysLysGln

1390 1410 1430

TCACTGCAGTGCTGATGTATTTCAAGTCTCTCACCCTCATCACTAGGAGATTCCTTTCCC

1450 1470 1490

CTGTAGGGTAGAGGGGTGGGGACAGAAACAACTTTCTCCTACTCTTCCTTCCTAATAGGC

1510 1530 1550

ATCTCCAGGCTGCCTGGTCACTGCCCCTCTCTCAGTGTCAATAGATGAAAGTACATTGGG

1570 1590 1610

AGTCTGTAGGAAACCCAACCTTCTTGTCATTGAAATTTGGCAAAGCTGACTTTGGGAAAG

1630 1650 1670

AGGGACCAGAACTTCCCCTCCCTTCCCCTTTTCCCAACCTGGACTTGTTTTAAACTTGCC

1690 1710 1730

TGTTCAGAGCACTCATTCCTTCCCACCCCCAGTCCTGTCCTATCACTCTAATTCGGATTT

1750 1770 1790

GCCATAGCCTTGAGGTTATGTCCTTTTCCATTAAGTACATGTGCCAGGAAACAGCGAGAG

1810 1830 1850

AGAGAAAGTAAACGGCAGTAATGCTTCTCCTATTTCTCCAAAGCCTTGTGTGAACTAGCA

1870 1890 1910

AAGAGAAGAAAATCAAATATATAACCAATAGTGAAATGCCACAGGTTTGTCCACTGTCAG

1930 1950 1970

GGTTGTCTACCTGTAGGATCAGGGTCTAAGCACCTTGGTGCTTAGCTAGAATACCACCTA

1990 2010 2030

ATCCTTCTGGCAAGCCTGTCTTCAGAGAACCCACTAGAAGCAACTAGGAAAAATCACTTG

2050 2070 2090

CCAAAATCCAAGGCAATTCCTGATGGAAAATGCAAAAGCACATATATGTTTTAATATCTT

2110 2130 2150

TATGGGCTCTGTTCAAGGCAGTGCTGAGAGGGAGGGGTTATAGCTTCAGGAGGGAACCAG

2170 2190 2210

CTTCTGATAAACACAATCTGCTAGGAACTTGGGAAAGGAATCAGAGAGCTGCCCTTCAGC

2230 2250 2270

GATTATTTAAATTGTTAAAGAATACACAATTTGGGGTATTGGGATTTTTCTCCTTTTCTC

2290 2310 2330

TGAGACATTCCACCATTTTAATTTTTGTAACTGCTTATTTATGTGAAAAGGGTTATTTTT

2350 2370 2390

ACTTAGCTTAGCTATGTCAGCCAATCCGATTGCCTTAGGTGAAAGAAACCACCGAAATCC

2410 2430 2450

CTCAGGTCCCTTGGTCAGGAGCCTCTCAAGATTTTTTTTGTCAGAGGCTCCAAATAGAAA

2470 2490 2510

ATAAGAAAAGGTTTTCTTCATTCATGGCTAGAGCTAGATTTAACTCAGTTTCTAGGCACC

2530 2550 2570

TCAGACCAATCATCAACTACCATTCTATTCCATGTTTGCACCTGTGCATTTTCTGTTTGC

2590 2610 2630

CCCCATTCACTTTGTCAGGAAACCTTGGCCTCTGCTAAGGTGTATTTGGTCCTTGAGAAG

2650 2670 2690

TGGGAGCACCCTACAGGGACACTATCACTCATGCTGGTGGCATTGTTTACAGCTAGAAAG

2710 2730 2750

CTGCACTGGTGCTAATGCCCCTTGGGAAATGGGGCTGTGAGGAGGAGGATTATAACTTAG

2770 2790 2810

GCCTAGCCTCTTTTAACAGCCTCTGAAATTTATCTTTTCTTCTATGGGGTCTATAAATGT

2830 2850 2870

ATCTTATAATAAAAAGGAAGGACAGGAGGAAGACAGGCAAATGTACTTCTCACCCAGTCT

2890 2910 2930

TCTACACAGATGGAATCTCTTTGGGGCTAAGAGAAAGGTTTTATTCTATATTGCTTACCT

2950 2970 2990

GATCTCATGTTAGGCCTAAGAGGCTTTCTCCAGGAGGATTAGCTTGGAGTTCTCTATACT

3010 3030 3050

CAGGTACCTCTTTCAGGGTTTTCTAACCCTGACACGGACTGTGCATACTTTCCCTCATCC

3070 3090 3110

ATGCTGTGCTGTGTTATTTAATTTTTCCTGGCTAAGATCATGTCTGAATTATGTATGAAA

3130 3150 3170

ATTATTCTATGTTTTTATAATAAAAATAATATATCAGACATCGAAAAAAAAAAA

SEQUENCE AND TRANSLATION OF cDNA OF TM-3

```
          10                  30                  50
CAGCCGTGCTCGAAGCGTTCCTGGAGCCCAAGCTCTCCTCCACAGGTGAAGACAGGGCCA

          70                  90                 110
GCAGGAGACACCATGGGGCACCTCTCAGCCCCACTTCACAGAGTGCGTGTACCCTGGCAG
             MetGlyHisLeuSerAlaProLeuHisArgValArgValProTrpGln

         130                 150                 170
GGGCTTCTGCTCACAGCCTCACTTCTAACCTTCTGGAACCCGCCCACCACTGCCCAGCTC
GlyLeuLeuLeuThrAlaSerLeuLeuThrPheTrpAsnProProThrThrAlaGlnLeu

         190                 210                 230
ACTACTGAATCCATGCCATTCAATGTTGCAGAGGGGAAGGAGGTTCTTCTCCTTGTCCAC
ThrThrGluSerMetProPheAsnValAlaGluGlyLysGluValLeuLeuLeuValHis

         250                 270                 290
AATCTGCCCCAGCAACTTTTTGGCTACAGCTGGTACAAAGGGGAAAGAGTGGATGGCAAC
AsnLeuProGlnGlnLeuPheGlyTyrSerTrpTyrLysGlyGluArgValAspGlyAsn

         310                 330                 350
CGTCAAATTGTAGGATATGCAATAGGAACTCAACAAGCTACCCCAGGGCCCGCAAACAGC
ArgGlnIleValGlyTyrAlaIleGlyThrGlnGlnAlaThrProGlyProAlaAsnSer

         370                 390                 410
GGTCGAGAGACAATATACCCCAATGCATCCCTGCTGATCCAGAACGTCACCCAGAATGAC
GlyArgGluThrIleTyrProAsnAlaSerLeuLeuIleGlnAsnValThrGlnAsnAsp
```

```
          430                 450                 470
          .                   .                   .
ACAGGATTCTACACCCTACAAGTCATAAAGTCAGATCTTGTGAATGAAGAAGCAACTGGA
ThrGlyPheTyrThrLeuGlnValIleLysSerAspLeuValAsnGluGluAlaThrGly

          490                 510                 530
          .                   .                   .
CAGTTCCATGTATACCCGGAGCTGCCCAAGCCCTCCATCTCCAGCAACAACTCCAACCCT
GlnPheHisValTyrProGluLeuProLysProSerIleSerSerAsnAsnSerAsnPro

          550                 570                 590
          .                   .                   .
GTGGAGGACAAGGATGCTGTGGCCTTCACCTGTGAACCTGAGACTCAGGACACAACCTAC
ValGluAspLysAspAlaValAlaPheThrCysGluProGluThrGlnAspThrThrTyr

          610                 630                 650
          .                   .                   .
CTGTGGTGGATAAACAATCAGAGCCTCCCGGTCAGTCCCAGGCTGCAGCTGTCCAATGGC
LeuTrpTrpIleAsnAsnGlnSerLeuProValSerProArgLeuGlnLeuSerAsnGly

          670                 690                 710
          .                   .                   .
AACAGGACCCTCACTCTACTCAGTGTCACAAGGAATGACACAGGACCCTATGAGTGTGAA
AsnArgThrLeuThrLeuLeuSerValThrArgAsnAspThrGlyProTyrGluCysGlu

          730                 750                 770
          .                   .                   .
ATACAGAACCCAGTGAGTGCGAACCGCAGTGACCCAGTCACCTTGAATGTCACCTATGGC
IleGlnAsnProValSerAlaAsnArgSerAspProValThrLeuAsnValThrTyrGly

          790                 810                 830
          .                   .                   .
CCGGACACCCCCACCATTTCCCCTTCAGACACCTATTACCGTCCAGGGGCAAACCTCAGC
ProAspThrProThrIleSerProSerAspThrTyrTyrArgProGlyAlaAsnLeuSer
```

```
          850                 870                 890
CTCTCCTGCTATGCAGCCTCTAACCCACCTGCACAGTACTCCTGGCTTATCAATGGAACA
LeuSerCysTyrAlaAlaSerAsnProProAlaGlnTyrSerTrpLeuIleAsnGlyThr

          910                 930                 950
TTCCAGCAAAGCACACAAGAGCTCTTTATCCCTAACATCACTGTGAATAATAGTGGATCC
PheGlnGlnSerThrGlnGluLeuPheIleProAsnIleThrValAsnAsnSerGlySer

          970                 990                 1010
TATACCTGCCACGCCAATAACTCAGTCACTGGCTGCAACAGGACCACAGTCAAGACGATC
TyrThrCysHisAlaAsnAsnSerValThrGlyCysAsnArgThrThrValLysThrIle

          1030                1050                1070
ATAGTCACTGAGCTAAGTCCAGTAGTAGCAAAGCCCCAAATCAAAGCCAGCAAGACCACA
IleValThrGluLeuSerProValValAlaLysProGlnIleLysAlaSerLysThrThr

          1090                1110                1130
GTCACAGGAGATAAGGACTCTGTGAACCTGACCTGCTCCACAAATGACACTGGAATCTCC
ValThrGlyAspLysAspSerValAsnLeuThrCysSerThrAsnAspThrGlyIleSer

          1150                1170                1190
ATCCGTTGGTTCTTCAAAAACCAGAGTCTCCCGTCCTCGGAGAGGATGAAGCTGTCCCAG
IleArgTrpPhePheLysAsnGlnSerLeuProSerSerGluArgMetLysLeuSerGln

          1210                1230                1250
GGCAACACCACCCTCAGCATAAACCCTGTCAAGAGGGAGGATGCTGGGACGTATTGGTGT
GlyAsnThrThrLeuSerIleAsnProValLysArgGluAspAlaGlyThrTyrTrpCys
```

```
      1270                1290                1310
GAGGTCTTCAACCCAATCAGTAAGAACCAAAGCGACCCCATCATGCTGAACGTAAACTAT
GluValPheAsnProIleSerLysAsnGlnSerAspProIleMetLeuAsnValAsnTyr

      1330                1350                1370
AATGCTCTACCACAAGAAAATGGCCTCTCACCTGGGGCCATTGCTGGCATTGTGATTGGA
AsnAlaLeuProGlnGluAsnGlyLeuSerProGlyAlaIleAlaGlyIleValIleGly

      1390                1410                1430
GTAGTGGCCCTGGTTGCTCTGATAGCAGTAGCCCTGGCATGTTTTCTGCATTTCGGGAAG
ValValAlaLeuValAlaLeuIleAlaValAlaLeuAlaCysPheLeuHisPheGlyLys

      1450                1470                1490
ACCGGCAGCTCAGGACCACTCCAATGACCCACCTAACAAGATGAATGAAGTTACTTATTC
ThrGlySerSerGlyProLeuGln

      1510                1530                1550
TACCCTGAACTTTGAAGCCCAGCAACCCACACAACCAACTTCAGCCTCCCCATCCCTAAC

      1570                1590                1610
AGCCACAGAAATAATTTATTCAGAAGTAAAAAAGCAGTAATGAAACCTGAAAAAAAAAAA

      1630
AAAAAAAAAA
```

SEQUENCE AND TRANSLATION OF cDNA OF TM-4

```
          10                  30                  50
          .                   .                   .
CAGCCGTGCTCGAAGCGTTCCTGGAGCCCAAGCTCTCCTCCACAGGTGAAGACAGGGCCA

          70                  90                  110
          .                   .                   .
GCAGGAGACACCATGGGGCACCTCTCAGCCCCACTTCACAGAGTGCGTGTACCCTGGCAG
             MetGlyHisLeuSerAlaProLeuHisArgValArgValProTrpGln

        130                 150                 170
         .                   .                   .
GGGCTTCTGCTCACAGCCTCACTTCTAACCTTCTGGAACCCGCCCACCACTGCCCAGCTC
GlyLeuLeuLeuThrAlaSerLeuLeuThrPheTrpAsnProProThrThrAlaGlnLeu

        190                 210                 230
         .                   .                   .
ACTACTGAATCCATGCCATTCAATGTTGCAGAGGGGAAGGAGGTTCTTCTCCTTGTCCAC
ThrThrGluSerMetProPheAsnValAlaGluGlyLysGluValLeuLeuLeuValHis

        250                 270                 290
         .                   .                   .
AATCTGCCCCAGCAACTTTTTGGCTACAGCTGGTACAAAGGGGAAAGAGTGGATGGCAAC
AsnLeuProGlnGlnLeuPheGlyTyrSerTrpTyrLysGlyGluArgValAspGlyAsn

        310                 330                 350
         .                   .                   .
CGTCAAATTGTAGGATATGCAATAGGAACTCAACAAGCTACCCCAGGGCCCGCAAACAGC
ArgGlnIleValGlyTyrAlaIleGlyThrGlnGlnAlaThrProGlyProAlaAsnSer

        370                 390                 410
         .                   .                   .
GGTCGAGAGACAATATACCCCAATGCATCCCTGCTGATCCAGAACGTCACCCAGAATGAC
GlyArgGluThrIleTyrProAsnAlaSerLeuLeuIleGlnAsnValThrGlnAsnAsp

        430                 450                 470
         .                   .                   .
ACAGGATTCTACACCCTACAAGTCATAAAGTCAGATCTTGTGAATGAAGAAGCAACTGGA
ThrGlyPheTyrThrLeuGlnValIleLysSerAspLeuValAsnGluGluAlaThrGly
```

```
          490                 510                 530
CAGTTCCATGTATACCCGGAGCTGCCCAAGCCCTCCATCTCCAGCAACAACTCCAACCCT
GlnPheHisValTyrProGluLeuProLysProSerIleSerSerAsnAsnSerAsnPro

          550                 570                 590
GTGGAGGACAAGGATGCTGTGGCCTTCACCTGTGAACCTGAGACTCAGGACACAACCTAC
ValGluAspLysAspAlaValAlaPheThrCysGluProGluThrGlnAspThrThrTyr

          610                 630                 650
CTGTGGTGGATAAACAATCAGAGCCTCCCGGTCAGTCCCAGGCTGCAGCTGTCCAATGGC
LeuTrpTrpIleAsnAsnGlnSerLeuProValSerProArgLeuGlnLeuSerAsnGly

          670                 690                 710
AACAGGACCCTCACTCTACTCAGTGTCACAAGGAATGACACAGGACCCTATGAGTGTGAA
AsnArgThrLeuThrLeuLeuSerValThrArgAsnAspThrGlyProTyrGluCysGlu

          730                 750                 770
ATACAGAACCCAGTGAGTGCGAACCGCAGTGACCCAGTCACCTTGAATGTCACCTATGGC
IleGlnAsnProValSerAlaAsnArgSerAspProValThrLeuAsnValThrTyrGly

          790                 810                 830
CCGGACACCCCCACCATTTCCCCTTCAGACACCTATTACCGTCCAGGGGCAAACCTCAGC
ProAspThrProThrIleSerProSerAspThrTyrTyrArgProGlyAlaAsnLeuSer

          850                 870                 890
CTCTCCTGCTATGCAGCCTCTAACCCACCTGCACAGTACTCCTGGCTTATCAATGGAACA
LeuSerCysTyrAlaAlaSerAsnProProAlaGlnTyrSerTrpLeuIleAsnGlyThr

          910                 930                 950
TTCCAGCAAAGCACACAAGAGCTCTTTATCCCTAACATCACTGTGAATAATAGTGGATCC
PheGlnGlnSerThrGlnGluLeuPheIleProAsnIleThrValAsnAsnSerGlySer

          970                 990                 1010
TATACCTGCCACGCCAATAACTCAGTCACTGGCTGCAACAGGACCACAGTCAAGACGATC
TyrThrCysHisAlaAsnAsnSerValThrGlyCysAsnArgThrThrValLysThrIle
```

```
      1030                1050                1070
ATAGTCACTGATAATGCTCTACCACAAGAAAATGGCCTCTCACCTGGGGCCATTGCTGGC
IleValThrAspAsnAlaLeuProGlnGluAsnGlyLeuSerProGlyAlaIleAlaGly

      1090                1110                1130
ATTGTGATTGGAGTAGTGGCCCTGGTTGCTCTGATAGCAGTAGCCCTGGCATGTTTTCTG
IleValIleGlyValValAlaLeuValAlaLeuIleAlaValAlaLeuAlaCysPheLeu

      1150                1170                1190
CATTTCGGGAAGACCGGCAGCTCAGGACCACTCCAATGACCCACCTAACAAGATGAATGA
HisPheGlyLysThrGlySerSerGlyProLeuGln

      1210                1230                1250
AGTTACTTATTCTACCCTGAACTTTGAAGCCCAGCAACCCACACAACCAACTTCAGCCTC

      1270                1290                1310
CCCATCCCTAACAGCCACAGAAATAATTTATTCAGAAGTAAAAAAAGCAGTAATGAAACCT

      1330
GAAAAAAAAAAAAAAAAAA
```

The present invention is also directed to a replicable recombinant cloning vehicle ("vector") having an insert comprising a nucleic acid, e.g., DNA, which comprises a base sequence which codes for a CEA peptide or a base sequence hybridizable therewith.

This invention also relates to a cell that is transformed/transfected, infected or injected with the above described replicable recombinant cloning vehicle or nucleic acid hybridizable with the aforementioned cDNA. Thus the invention also concerns the transfection of cells using free nucleic acid, without the use of a cloning vehicle.

Still further, the present invention concerns a polypeptide expressed by the above described transfected, infected or injected cell, which polypeptide exhibits immunological cross-reactivity with a CEA, as well as labelled forms of the polypeptide. The invention also relates to polypeptides having an amino acid sequence, i.e., synthetic peptides, or the expression product of a cell that is transfected, injected, infected with the above described replicable recombinant cloning vehicles, as well as labelled forms thereof. Stated otherwise, the present invention concerns a synthetic peptide having an amino acid sequence corresponding to the entire amino acid sequence or a portion thereof having no less than five amino acids of the aforesaid expression product.

The invention further relates to an antibody preparation specific for the above described polypeptide.

Another aspect of the invention concerns an immunoassay method for detecting CEA or a functional equivalent thereof in a test sample comprising

(a) contacting the sample with the above described antibody preparation, and

(b) determining binding thereof to CEA in the sample.

The invention also is directed to a nucleic acid hybridization method for detecting a CEA or a related nucleic acid (DNA or RNA) sample in a test sample comprising

(a) contacting the test sample with a nucleic acid probe comprising a nucleic acid, which comprises a base sequence which codes for a CEA peptide sequence or a base sequence that is hybridizable therewith, and

(b) determining the formation of the resultant hybridized probe.

The present invention also concerns a method for detecting the presence of carcinoembryonic antigen or a functional equivalent thereof in an animal or human patient in vivo comprising

a) introducing into said patient a labeled (e.g., a radio-opaque material that can be detected by X-rays, radiolabeled or labeled with paramagnetic materials that can be detected by NMR) antibody preparation according to the present invention and

b) detecting the presence of such antibody preparation in the patient by detecting the label.

In another aspect, the present invention relates to the use of an antibody preparation according to the present invention for therapeutic purposes, namely, attaching to an antibody preparation radionuclides, toxins or other biological effectors to form a complex and introducing an effective amount of such complex into an animal or human patient, e.g., by injection or orally. The antibody complex would attach to CEA in a patient and the radionuclide, toxin or other biological effector would serve to destroy the CEA expressing cell.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 is a schematic representation of the transmembrane CEA's

DETAILED DESCRIPTION OF THE INVENTION

In parent applications, applicants described the following CEA's:

		ATCC No.
CEA-(a)	partial CEA (pcLV7)	
CEA-(b)	full coding CEA (pc 15LV7)	67709
CEA-(c)	TM-1 (FL-CEA; pc 19-22)	67710
CEA-(d)	NCA (pcBT 20)	67711

In the present application, applicants described the following CEA's:

		ATCC No.
CEA-(e)	TM-2 (pc E22)	67712
CEA-(f)	TM-3 (pc HT-6)	67708
CEA-(g)	TM-4.	

ATCC Nos. 67708, 67709, 67710, 67711 and 67712 were all deposited with the American Type Culture Collection on May 25, 1988.

The sequences for CEA-(a), CEA-(b), CEA-(c) and CEA-(d) are given hereinbelow:

CEA-(a):

GG GGT TTA CAC AAC CAC CAC CCC ATC AAA CCC TTC ATC ACC AGC AAC AAC TCC AAC CCC GTG

GAG GAT GAG GAT GCT GTA GCC TTA ACC TGT GAA CCT GAG ATT CAG AAC ACA ACC TAC CTG

TGG TGG GTA AAT AAT CAG AGC CTC CCG GTC AGT CCC AGG CTG CAG CTG TCC AAT GAC AAC

AGG ACC CTC ACT CTA CTC AGT GTC ACA AGG AAT GAT GTA GGA CCC TAT GAG TGT GGA ATC

CAG AAC GAA TTA AGT GTT GAC CAC AGC GAC CCA GTC ACC CAG CGA TTC CTC TAT GGC CCA

GAC GAC CCC ACC ATT TCC CCC TCA TAC ACC TAT TAC CGT CCA GGG GTG GAA CCT CAG CCT

CTC TCC CAT GCA GCC TCT AAC CCA CCT GCA CAG TAT TCT TGG CTG ATT GAT GGG ACC GTC

CAG CAA CAC ACA CAA GAG CTC TTT ATC TCC AAC ATC ACT GAG AAG AAC AGC GGA CTC TAT

ACC TGC CAG GCC AAT AAC TCA GCC AGT GGC ACA GCA GGA CTA CAG TCA AGA CAA TCA CAG

TCT CTG CGG ATG CCC AAG CCC TCC ATC TCC AGC AAC AAC TCC AAA CCC GTG GAG GAC AAG

GAT CGC TGT GGC CTT CAC TGT GAA CCT GAG GCT CAG AAC ACA ACC TAC CTG TGG TGG GTA

AAT GGT CAG AGC CTC CCA GTC AGT CCC AGG CTG CAG CTG TCC AAT GGC AAC AGG ACC CTC

ACT CTA TTC AAT GTC ACA AGA AAT GAC GCA AGA GCC TAT GTA TGT GGA ATC CAG AAC TCA

GTG AGT GCA AAC CGC AGT GAC CCA GTC ACC CTG GAT GTC CTC TAT GGG CCG GAC ACC CCC

ATC ATT TCC CCC CCC CC

(b)

```
                10          20          30          40          50
                *           *           *           *           *
C ACC ATG GAG TCT CCC TCG GCC CCT CTC CAC AGA TGG TGC ATC CCC TGG CAG AGG CTC
      Met Glu Ser Pro Ser Ala Pro Leu His Arg Trp Cys Ile Pro Trp Gln Arg Leu
```

```
          60          70          80          90          100         110
          *           *           *           *           *           *
CTG CTC ACA GCC TCA CTT CTA ACC TTC TGG AAC CCG CCC ACC ACT GCC AAG CTC ACT
Leu Leu Thr Ala Ser Leu Leu Thr Phe Trp Asn Pro Pro Thr Thr Ala Lys Leu Thr
                                                                 1   2   3

      120         130         140         150         160         170
       *           *           *           *           *           *
ATT GAA TCC ACG CCG TTC AAT GTC GCA GAG GGG AAG GAG GTG CTT CTA CTT GTC CAC
Ile Glu Ser Thr Pro Phe Asn Val Ala Glu Gly Lys Glu Val Leu Leu Leu Val His
  4   5   6   7   8   9  10  11  12  13  14  15  16  17  18  19  20  21  22

            180         190             200         210         220
             *           *               *           *           *
AAT CTG CCC CAG CAT CTT TTT GGC TAC AGC TGG TAC AAA GGT GAA AGA GTG GAT GGC
Asn Leu Pro Gln His Leu Phe Gly Tyr Ser Trp Tyr Lys Gly Glu Arg Val Asp Gly
 23  24  25  26  27  28  29  30  31  32  33  34  35  36  37  38  39  40  41

230         240         250         260         270         280
 *           *           *           *           *           *
AAC CGT CAA ATT ATA GGA TAT GTA ATA GGA ACT CAA CAA GCT ACC CCA GGG CCC GCA
Asn Arg Gln Ile Ile Gly Tyr Val Ile Gly Thr Gln Gln Ala Thr Pro Gly Pro Ala
 42  43  44  45  46  47  48  49  50  51  52  53  54  55  56  57  58  59  60

   290         300         310         320         330         340
    *           *           *           *           *           *
TAC AGT GGT CGA GAG ATA ATA TAC CCC AAT GCA TCC CTG CTG ATC CAG AAC ATC ATC
Tyr Ser Gly Arg Glu Ile Ile Tyr Pro Asn Ala Ser Leu Leu Ile Gln Asn Ile Ile
 61  62  63  64  65  66  67  68  69  70  71  72  73  74  75  76  77  78  79

        350         360         370         380         390         400
         *           *           *           *           *           *
CAG AAT GAC ACA GGA TTC TAC ACC CTA CAC GTC ATA AAG TCA GAT CTT GTG AAT GAA
Gln Asn Asp Thr Gly Phe Tyr Thr Leu His Val Ile Lys Ser Asp Leu Val Asn Glu
 80  81  82  83  84  85  86  87  88  89  90  91  92  93  94  95  96  97  98

            410             420         430         440             450
             *               *           *           *               *
GAA GCA ACT GGC CAG TTC CGG GTA TAC CCG GAG CTG CCC AAG CCC TCC ATC TCC AGC
Glu Ala Thr Gly Gln Phe Arg Val Tyr Pro Glu Leu Pro Lys Pro Ser Ile Ser Ser
 99 101 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117

 460         470         480         490         500         510
  *           *           *           *           *           *
AAC AAC TCC AAA CCC GTG GAG GAC AAG GAT GCT GTG GCC TTC ACC TGT GAA CCT GAG
Asn Asn Ser Lys Pro Val Glu Asp Lys Asp Ala Val Ala Phe Thr Cys Glu Pro Glu
118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136
```

```
      520         530         540         550         560         570
       *           *           *           *           *           *
ACT CAG GAC GCA ACC TAC CTG TGG TGG GTA AAC AAT CAG AGC CTC CCG GTC AGT CCC
Thr Gln Asp Ala Thr Tyr Leu Trp Trp Val Asn Asn Gln Ser Leu Pro Val Ser Pro
137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155

         580         590         600         610         620
          *           *           *           *           *
AGG CTG CAG CTG TCC AAT GGC AAC AGG ACC CTC ACT CTA TTC AAT GTC ACA AGA AAT
Arg Leu Gln Leu Ser Asn Gly Asn Arg Thr Leu Thr Leu Phe Asn Val Thr Arg Asn
156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174

630         640         650         660         670         680
 *           *           *           *           *           *
GAA CAA GCA AGC TAC AAA TGT GAA ACC CAG AAC CCA GTG AGT GCC AGG CGC AGT GAT
Glu Gln Ala Ser Tyr Lys Cys Glu Thr Gln Asn Pro Val Ser Ala Arg Arg Ser Asp
175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193

      690         700         710         720         730         740
       *           *           *           *           *           *
TCA GTC ATC CTG AAT GTC CTC TAT GGC CCG GAT GCC CCC ACC ATT TCC CCT CTA AAC
Ser Val Ile Leu Asn Val Leu Tyr Gly Pro Asp Ala Pro Thr Ile Ser Pro Leu Asn
194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212

         750         760         770         780         790
          *           *           *           *           *
ACA TCT TAC AGA TCA GGG GAA AAT CTG AAC CTC TCC TGC CAC GCA GCC TCT AAC CCA
Thr Ser Tyr Arg Ser Gly Glu Asn Leu Asn Leu Ser Cys His Ala Ala Ser Asn Pro
213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231

800         810         820         830         840         850
 *           *           *           *           *           *
CCT GCA CAG TAC TCT TGG TTT GTC AAT GGG ACT TTC CAG CAA TCC ACC CAA GAG CTC
Pro Ala Gln Tyr Ser Trp Phe Val Asn Gly Thr Phe Gln Gln Ser Thr Gln Glu Leu
232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250

      860         870         880         890         900         910
       *           *           *           *           *           *
TTT ATC CCC AAC ATC ACT GTG AAT AAT AGT GGA TCC TAT ACG TGC CAA GCC CAT AAC
Phe Ile Pro Asn Ile Thr Val Asn Asn Ser Gly Ser Tyr Thr Cys Gln Ala His Asn
251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269

         920         930         940         950         960         970
          *           *           *           *           *           *
TCA GAC ACT GGC CTC AAT AGG ACC ACA GTC ACG ACG ATC ACA GTC TAT GCA GAG CCA
Ser Asp Thr Gly Leu Asn Arg Thr Thr Val Thr Thr Ile Thr Val Tyr Ala Glu Pro
270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288
```

```
                980         990         1000        1010        1020
                 *           *           *           *           *
CCC AAA CCC TTC ATC ACC AGC AAC AAC TCC AAC CCC GTG GAG GAT GAG GAT GCT GTA
Pro Lys Pro Phe Ile Thr Ser Asn Asn Ser Asn Pro Val Glu Asp Glu Asp Ala Val
289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307

1030        1040        1050        1060        1070        1080
 *           *           *           *           *           *
GCC TTA ACC TGT GAA CCT GAG ATT CAG AAC ACA ACC TAC CTG TGG TGG GTA AAT AAT
Ala Leu Thr Cys Glu Pro Glu Ile Gln Asn Thr Thr Tyr Leu Trp Trp Val Asn Asn
308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326

1090        1100        1110        1120        1130        1140
 *           *           *           *           *           *
CAG AGC CTC CCG GTC AGT CCC AGG CTG CAG CTG TCC AAT GAC AAC AGG ACC CTC ACT
Gln Ser Leu Pro Val Ser Pro Arg Leu Gln Leu Ser Asn Asp Asn Arg Thr Leu Thr
327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345

            1150        1160        1170        1180        1190
             *           *           *           *           *
CTA CTC AGT GTC ACA AGG AAT GAT GTA GGA CCC TAT GAG TGT GGA ATC CAG AAC GAA
Leu Leu Ser Val Thr Arg Asn Asp Val Gly Pro Tyr Glu Cys Gly Ile Gln Asn Glu
346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364

1200        1210        1220        1230        1240        1250
 *           *           *           *           *           *
TTA AGT GTT GAC CAC AGC GAC CCA GTC ATC CTG AAT GTC CTC TAT GGC CCA GAC GAC
Leu Ser Val Asp His Ser Asp Pro Val Ile Leu Asn Val Leu Tyr Gly Pro Asp Asp
365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383

   1260        1270        1280        1290        1300        1310
    *           *           *           *           *           *
CCC ACC ATT TCC CCC TCA TAC ACC TAT TAC CGT CCA GGG GTG AAC CTC AGC CTC TCC
Pro Thr Ile Ser Pro Ser Tyr Thr Tyr Tyr Arg Pro Gly Val Asn Leu Ser Leu Ser
384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402

         1320        1330        1340        1350        1360
          *           *           *           *           *
TGC CAT GCA GCC TCT AAC CCA CCT GCA CAG TAT TCT TGG CTG ATT GAT GGG AAC ATC
Cys His Ala Ala Ser Asn Pro Pro Ala Gln Tyr Ser Trp Leu Ile Asp Gly Asn Ile
403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421

1370        1380        1390        1400        1410        1420
 *           *           *           *           *           *
CAG CAA CAC ACA CAA GAG CTC TTT ATC TCC AAC ATC ACT GAG AAG AAC AGC GGA CTC
Gln Gln His Thr Gln Glu Leu Phe Ile Ser Asn Ile Thr Glu Lys Asn Ser Gly Leu
422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440
```

```
      1430        1440        1450        1460        1470        1480
       *           *           *           *           *           *
TAT ACC TGC CAG GCC AAT AAC TCA GCC AGT GGC CAC AGC AGG ACT ACA GTC AAG ACA
Tyr Thr Cys Gln Ala Asn Asn Ser Ala Ser Gly His Ser Arg Thr Thr Val Lys Thr
441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459

      1490        1500        1510        1520        1530        1540
       *           *           *           *           *           *
ATC ACA GTC TCT GCG GAC GTG CCC AAG CCC TCC ATC TCC AGC AAC AAC TCC AAA CCC
Ile Thr Val Ser Ala Asp Val Pro Lys Pro Ser Ile Ser Ser Asn Asn Ser Lys Pro
460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478

      1550        1560        1570        1580        1590
       *           *           *           *           *
GTG GAG GAC AAG GAT GCT GTG GCC TTC ACC TGT GAA CCT GAG GCT CAG AAC ACA ACC
Val Glu Asp Lys Asp Ala Val Ala Phe Thr Cys Glu Pro Glu Ala Gln Asn Thr Thr
479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497

      1600        1610        1620        1630        1640        1650
       *           *           *           *           *           *
TAC CTG TGG TGG GTA AAT GGT CAG AGC CTC CCA GTC AGT CCC AGG CTG CAG CTG TCC
Tyr Leu Trp Trp Val Asn Gly Gln Ser Leu Pro Val Ser Pro Arg Leu Gln Leu Ser
498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516

      1660        1670        1680        1690        1700        1710
       *           *           *           *           *           *
AAT GGC AAC AGG ACC CTC ACT CTA TTC AAT GTC ACA AGA AAT GAC GCA AGA GCC TAT
Asn Gly Asn Arg Thr Leu Thr Leu Phe Asn Val Thr Arg Asn Asp Ala Arg Ala Tyr
517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535

      1720        1730        1740        1750        1760
       *           *           *           *           *
GTA TGT GGA ATC CAG AAC TCA GTG AGT GCA AAC CGC AGT GAC CCA GTC ACC CTG GAT
Val Cys Gly Ile Gln Asn Ser Val Ser Ala Asn Arg Ser Asp Pro Val Thr Leu Asp
536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554

1770        1780        1790        1800        1810        1820
 *           *           *           *           *           *
GTC CTC TAT GGG CCG GAC ACC CCC ATC ATT TCC CCC CCA GAC TCG TCT TAC CTT TCG
Val Leu Tyr Gly Pro Asp Thr Pro Ile Ile Ser Pro Pro Asp Ser Ser Tyr Leu Ser
555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573

      1830        1840        1850        1860        1870        1880
       *           *           *           *           *           *
GGA GCG AAC CTC AAC CTC TCC TGC CAC TCG GCC TCT AAC CCA TCC CCG CAG TAT TCT
Gly Ala Asn Leu Asn Leu Ser Cys His Ser Ala Ser Asn Pro Ser Pro Gln Tyr Ser
574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592
```

```
          1890        1900        1910        1920        1930
           *           *           *           *           *
TGG CGT ATC AAT GGG ATA CCG CAG CAA CAC ACA CAA GTT CTC TTT ATC GCC AAA ATC
Trp Arg Ile Asn Gly Ile Pro Gln Gln His Thr Gln Val Leu Phe Ile Ala Lys Ile
593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611

1940        1950        1960        1970        1980        1990
 *           *           *           *           *           *
ACG CCA AAT AAT AAC GGG ACC TAT GCC TGT TTT GTC TCT AAC TTG GCT ACT GGC CGC
Thr Pro Asn Asn Asn Gly Thr Tyr Ala Cys Phe Val Ser Asn Leu Ala Thr Gly Arg
612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630

2000        2010        2020        2030        2040        2050
 *           *           *           *           *           *
AAT AAT TCC ATA GTC AAG AGC ATC ACA GTC TCT GCA TCT GGA ACT TCT CCT GGT CTC
Asn Asn Ser Ile Val Lys Ser Ile Thr Val Ser Ala Ser Gly Thr Ser Pro Gly Leu
631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649

          2060        2070        2080        2090        2100        2110
           *           *           *           *           *           *
TCA GCT GGG GCC ACT GTC GGC ATC ATG ATT GGA GTG CTG GTT GGG GTT GCT CTG ATA
Ser Ala Gly Ala Thr Val Gly Ile Met Ile Gly Val Leu Val Gly Val Ala Leu Ile
650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668

                    2120        2130        2140        2150        2160
                     *           *           *           *           *
TAG CAG CCC TGG TGT AGT TTC TTC ATT TCA GGA AGA CTG ACA GTT GTT TTG CTT CTT

2170        2180        2190        2200        2210        2220
 *           *           *           *           *           *
CCT TAA AGC ATT TGC AAC AGC TAC AGT CTA AAA TTG CTT CTT TAC CAA GGA TAT TTA

    2230        2240        2250        2260        2270        2280
     *           *           *           *           *           *
CAG AAA ATA CTC TGA CCA GAG ATC GAG ACC ATC CTA GCC AAC ATC GTG AAA CCC CAT

          2290        2300        2310        2320        2330
           *           *           *           *           *
CTC TAC TAA AAA TAC AAA AAT GAG CTG GGC TTG GTG GCG CGC ACC TGT AGT CCC AGT

2340        2350        2360        2370        2380        2390
 *           *           *           *           *           *
TAC TCG GGA GGC TGA GGC AGG AGA ATC GCT TGA ACC CGG GAG GTG GAG ATT GCA GTG
```

2400 2410 2420 2430 2440 2450

AGC CCA GAT CGC ACC ACT GCA CTC CAG TCT GGC AAC AGA GCA AGA CTC CAT CTC AAA

2460 2470 2480 2490 2500

AAG AAA AGA AAA GAA GAC TCT GAC CTG TAC TCT TGA ATA CAA GTT TCT GAT ACC ACT

2510 2520 2530 2540 2550 2560

GCA CTG TCT GAG AAT TTC CAA AAC TTT AAT GAA CTA ACT GAC AGC TTC ATG AAA CTG

2570 2580 2590 2600 2610 2620

TCC ACC AAG ATC AAG CAG AGA AAA TAA TTA ATT TCA TGG GGA CTA AAT GAA CTA ATG

2630 2640 2650 2660 2670 2680

AGG ATA ATA TTT TCA TAA TTT TTT ATT TGA AAT TTT GCT GAT TCT TTA AAT GTC TTG

2690 2700 2710 2720 2730

TTT CCC AGA TTT CAG GAA ACT TTT TTT CTT TTA AGC TAT CCA CTC TTA CAG CAA TTT

2740 2750 2760 2770 2780 2790

GAT AAA ATA TAC TTT TGT GAA CAA AAA TTG AGA CAT TTA CAT TTT ATC CCT ATG TGG

2800 2810 2820 2830

TCG CTC CAG ACT TGG GAA ACT ATT CAT GAA TAT TTA TAT TGT ATG

CEA-(c):

```
          10                  30                  50
CAGCCGTGCTCGAAGCGTTCCTGGAGCCCAAGCTCTCCTCCACAGGTGAAGACAGGGCCA

          70                  90                 110
GCAGGAGACACCATGGGGCACCTCTCAGCCCCACTTCACAGAGTGCGTGTACCCTGGCAG
             MetGlyHisLeuSerAlaProLeuHisArgValArgValProTrpGln

         130                 150                 170
GGGCTTCTGCTCACAGCCTCACTTCTAACCTTCTGGAACCCGCCCACCACTGCCCAGCTC
GlyLeuLeuLeuThrAlaSerLeuLeuThrPheTrpAsnProProThrThrAlaGlnLeu

         190                 210                 230
ACTACTGAATCCATGCCATTCAATGTTGCAGAGGGGAAGGAGGTTCTTCTCCTTGTCCAC
ThrThrGluSerMetProPheAsnValAlaGluGlyLysGluValLeuLeuLeuValHis

         250                 270                 290
AATCTGCCCCAGCAACTTTTTGGCTACAGCTGGTACAAAGGGGAAAGAGTGGATGGCAAC
AsnLeuProGlnGlnLeuPheGlyTyrSerTrpTyrLysGlyGluArgValAspGlyAsn

         310                 330                 350
CGTCAAATTGTAGGATATGCAATAGGAACTCAACAAGCTACCCCAGGGCCCGCAAACAGC
ArgGlnIleValGlyTyrAlaIleGlyThrGlnGlnAlaThrProGlyProAlaAsnSer

         370                 390                 410
GGTCGAGAGACAATATACCCCAATGCATCCCTGCTGATCCAGAACGTCACCCAGAATGAC
GlyArgGluThrIleTyrProAsnAlaSerLeuLeuIleGlnAsnValThrGlnAsnAsp

         430                 450                 470
ACAGGATTCTACACCCTACAAGTCATAAAGTCAGATCTTGTGAATGAAGAAGCAACTGGA
ThrGlyPheTyrThrLeuGlnValIleLysSerAspLeuValAsnGluGluAlaThrGly

         490                 510                 530
```

```
CAGTTCCATGTATACCCGGAGCTGCCCAAGCCCTCCATCTCCAGCAACAACTCCAACCCT
GlnPheHisValTyrProGluLeuProLysProSerIleSerSerAsnAsnSerAsnPro

          550                 570                 590
GTGGAGGACAAGGATGCTGTGGCCTTCACCTGTGAACCTGAGACTCAGGACACAACCTAC
ValGluAspLysAspAlaValAlaPheThrCysGluProGluThrGlnAspThrThrTyr

          610                 630                 650
CTGTGGTGGATAAACAATCAGAGCCTCCCGGTCAGTCCCAGGCTGCAGCTGTCCAATGGC
LeuTrpTrpIleAsnAsnGlnSerLeuProValSerProArgLeuGlnLeuSerAsnGly

          670                 690                 710
AACAGGACCCTCACTCTACTCAGTGTCACAAGGAATGACACAGGACCCTATGAGTGTGAA
AsnArgThrLeuThrLeuLeuSerValThrArgAsnAspThrGlyProTyrGluCysGlu

          730                 750                 770
ATACAGAACCCAGTGAGTGCGAACCGCAGTGACCCAGTCACCTTGAATGTCACCTATGGC
IleGlnAsnProValSerAlaAsnArgSerAspProValThrLeuAsnValThrTyrGly

          790                 810                 830
CCGGACACCCCCACCATTTCCCCTTCAGACACCTATTACCGTCCAGGGGCAAACCTCAGC
ProAspThrProThrIleSerProSerAspThrTyrTyrArgProGlyAlaAsnLeuSer

          850                 870                 890
CTCTCCTGCTATGCAGCCTCTAACCCACCTGCACAGTACTCCTGGCTTATCAATGGAACA
LeuSerCysTyrAlaAlaSerAsnProProAlaGlnTyrSerTrpLeuIleAsnGlyThr

          910                 930                 950
TTCCAGCAAAGCACACAAGAGCTCTTTATCCCTAACATCACTGTGAATAATAGTGGATCC
PheGlnGlnSerThrGlnGluLeuPheIleProAsnIleThrValAsnAsnSerGlySer

          970                 990                 1010
TATACCTGCCACGCCAATAACTCAGTCACTGGCTGCAACAGGACCACAGTCAAGACGATC
TyrThrCysHisAlaAsnAsnSerValThrGlyCysAsnArgThrThrValLysThrIle

          1030                1050                1070
```

```
ATAGTCACTGAGCTAAGTCCAGTAGTAGCAAAGCCCCAAATCAAAGCCAGCAAGACCACA
IleValThrGluLeuSerProValValAlaLysProGlnIleLysAlaSerLysThrThr

      1090                1110                1130
GTCACAGGAGATAAGGACTCTGTGAACCTGACCTGCTCCACAAATGACACTGGAATCTCC
ValThrGlyAspLysAspSerValAsnLeuThrCysSerThrAsnAspThrGlyIleSer

      1150                1170                1190
ATCCGTTGGTTCTTCAAAAACCAGAGTCTCCCGTCCTCGGAGAGGATGAAGCTGTCCCAG
IleArgTrpPhePheLysAsnGlnSerLeuProSerSerGluArgMetLysLeuSerGln

      1210                1230                1250
GGCAACACCACCCTCAGCATAAACCCTGTCAAGAGGGAGGATGCTGGGACGTATTGGTGT
GlyAsnThrThrLeuSerIleAsnProValLysArgGluAspAlaGlyThrTyrTrpCys

      1270                1290                1310
GAGGTCTTCAACCCAATCAGTAAGAACCAAAGCGACCCCATCATGCTGAACGTAAACTAT
GluValPheAsnProIleSerLysAsnGlnSerAspProIleMetLeuAsnValAsnTyr

      1330                1350                1370
AATGCTCTACCACAAGAAAATGGCCTCTCACCTGGGGCCATTGCTGGCATTGTGATTGGA
AsnAlaLeuProGlnGluAsnGlyLeuSerProGlyAlaIleAlaGlyIleValIleGly

      1390                1410                1430
GTAGTGGCCCTGGTTGCTCTGATAGCAGTAGCCCTGGCATGTTTTCTGCATTTCGGGAAG
ValValAlaLeuValAlaLeuIleAlaValAlaLeuAlaCysPheLeuHisPheGlyLys

      1450                1470                1490
ACCGGCAGGGCAAGCGACCAGCGTGATCTCACAGAGCACAAACCCTCAGTCTCCAACCAC
ThrGlyArgAlaSerAspGlnArgAspLeuThrGluHisLysProSerValSerAsnHis

      1510                1530                1550
ACTCAGGACCACTCCAATGACCCACCTAACAAGATGAATGAAGTTACTTATTCTACCCTG
ThrGlnAspHisSerAsnAspProProAsnLysMetAsnGluValThrTyrSerThrLeu

      1570                1590                1610
```

```
AACTTTGAAGCCCAGCAACCCACACAACCAACTTCAGCCTCCCCATCCCTAACAGCCACA
AsnPheGluAlaGlnGlnProThrGlnProThrSerAlaSerProSerLeuThrAlaThr

      1630                1650                1670
GAAATAATTTATTCAGAAGTAAAAAAAGCAGTAATGAAACCTGTCCTGCTCACTGCAGTGC
GluIleIleTyrSerGluValLysLysGln

      1690                1710                1730
TGATGTATTTCAAGTCTCTCACCCTCATCACTAGGAGATTCCTTTCCCCTGTAGGGTAGA

      1750                1770                1790
GGGGTGGGGACAGAAACAACTTTCTCCTACTCTTCCTTCCTAATAGGCATCTCCAGGCTG

      1810                1830                1850
CCTGGTCACTGCCCCTCTCTCAGTGTCAATAGATGAAAGTACATTGGGAGTCTGTAGGAA

      1870                1890                1910
ACCCAACCTTCTTGTCATTGAAATTTGGCAAAGCTGACTTTGGGAAAGAGGGACCAGAAC

      1930                1950                1970
TTCCCCTCCCTTCCCCTTTTCCCAACCTGGACTTGTTTTAAACTTGCCTGTTCAGAGCAC

      1990                2010                2030
TCATTCCTTCCCACCCCCAGTCCTGTCCTATCACTCTAATTCGGATTTGCCATAGCCTTG

      2050                2070                2090
AGGTTATGTCCTTTTCCATTAAGTACATGTGCCAGGAAACAGCGAGAGAGAGAAAGTAAA

      2110                2130                2150
CGGCAGTAATGCTTCTCCTATTTCTCCAAAGCCTTGTGTGAACTAGCAAAGAGAAGAAAA

      2170                2190                2210
TCAAATATATAACCAATAGTGAAATGCCACAGGTTTGTCCACTGTCAGGGTTGTCTACCT
```

2230 2250 2270

GTAGGATCAGGGTCTAAGCACCTTGGTGCTTAGCTAGAATACCACCTAATCCTTCTGGCA

2290 2310 2330

AGCCTGTCTTCAGAGAACCCACTAGAAGCAACTAGGAAAAATCACTTGCCAAAATCCAAG

2350 2370 2390

GCAATTCCTGATGGAAAATGCAAAAGCACATATATGTTTTAATATCTTTATGGGCTCTGT

2410 2430 2450

TCAAGGCAGTGCTGAGAGGGAGGGGTTATAGCTTCAGGAGGGAACCAGCTTCTGATAAAC

2470 2490 2510

ACAATCTGCTAGGAACTTGGGAAAGGAATCAGAGAGCTGCCCTTCAGCGATTATTTAAAT

2530 2550 2570

TGTTAAAGAATACACAATTTGGGGTATTGGGATTTTTCTCCTTTTCTCTGAGACATTCCA

2590 2610 2630

CCATTTTAATTTTTGTAACTGCTTATTTATGTGAAAAGGGTTATTTTTACTTAGCTTAGC

2650 2670 2690

TATGTCAGCCAATCCGATTGCCTTAGGTGAAAGAAACCACCGAAATCCCTCAGGTCCCTT

2710 2730 2750

GGTCAGGAGCCTCTCAAGATTTTTTTTGTCAGAGGCTCCAAATAGAAAATAAGAAAAGGT

2770 2790 2810

TTTCTTCATTCATGGCTAGAGCTAGATTTAACTCAGTTTCTAGGCACCTCAGACCAATCA

2830 2850 2870

TCAACTACCATTCTATTCCATGTTTGCACCTGTGCATTTTCTGTTTGCCCCCATTCACTT

2890 2910 2930

TGTCAGGAAACCTTGGCCTCTGCTAAGGTGTATTTGGTCCTTGAGAAGTGGGAGCACCCT

2950 2970 2990

ACAGGGACACTATCACTCATGCTGGTGGCATTGTTTACAGCTAGAAAGCTGCACTGGTGC

3010 3030 3050

TAATGCCCCTTGGGAAATGGGGCTGTGAGGAGGAGGATTATAACTTAGGCCTAGCCTCTT

3070 3090 3110

TTAACAGCCTCTGAAATTTATCTTTTCTTCTATGGGGTCTATAAATGTATCTTATAATAA

3130 3150 3170

AAAGGAAGGACAGGAGGAAGACAGGCAAATGTACTTCTCACCCAGTCTTCTACACAGATG

3190 3210 3230

GAATCTCTTTGGGGCTAAGAGAAAGGTTTTATTCTATATTGCTTACCTGATCTCATGTTA

3250 3270 3290

GGCCTAAGAGGCTTTCTCCAGGAGGATTAGCTTGGAGTTCTCTATACTCAGGTACCTCTT

3310 3330 3350

TCAGGGTTTTCTAACCCTGACACGGACTGTGCATACTTTCCCTCATCCATGCTGTGCTGT

3370 3390 3410

GTTATTTAATTTTTCCTGGCTAAGATCATGTCTGAATTATGTATGAAAATTATTCTATGT

3430 3450

TTTTATAATAAAAATAATATATCAGACATCGAAAAAAAAAA

(d)

```
                10          20          30          40          50
                |           |           |           |           |
CC GGG GGA CAC GCA GGG CCA ACA GTC ACA GCA GCC CTG ACC AGA GCA TTC CTG GAG CTC

                60          70          80          90          100         110
                |           |           |           |           |           |
AAG CTC TCT ACA AAG AGG TGG ACA GAG AAG ACA GCA GAG ACC ATG GGA CCC CCC TCA
                                                            Met Gly Pro Pro Ser

                120         130         140         150         160         170
                |           |           |           |           |           |
GCC CCT CCC TGC AGA TTG CAT GTC CCC TGG AAG GAG GTC CTG CTC ACA GCC TCA CTT
Ala Pro Pro Cys Arg Leu His Val Pro Trp Lys Glu Val Leu Leu Thr Ala Ser Leu

                180         190         200         210         220         230
                |           |           |           |           |           |
CTA ACC TTC TGG AAC CCA CCC ACC ACT GCC AAG CTC ACT ATT GAA TCC ACG CCA TTC
Leu Thr Phe Trp Asn Pro Pro Thr Thr Ala Lys Leu Thr Ile Glu Ser Thr Pro Phe
                                                1  2  3  4  5  6  7  8  9
                240         250         260         270         280
                |           |           |           |           |
AAT GTC GCA GAG GGG AAG GAG GTT CTT CTA CTC GCC CAC AAC CTG CCC CAG AAT CGT
Asn Val Ala Glu Gly Lys Glu Val Leu Leu Leu Ala His Asn Leu Pro Gln Asn Arg
10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28

                290         300         310         320         330         340
                |           |           |           |           |           |
ATT GGT TAC AGC TGG TAC AAA GGC GAA AGA GTG GAT GGC AAC AGT CTA ATT GTA GGA
Ile Gly Tyr Ser Trp Tyr Lys Gly Glu Arg Val Asp Gly Asn Ser Leu Ile Val Gly
29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47

                350         360         370         380         390         400
                |           |           |           |           |           |
TAT GTA ATA GGA ACT CAA CAA GCT ACC CCA GGG CCC GCA TAC AGT GGT CGA GAG ACA
Tyr Val Ile Gly Thr Gln Gln Ala Thr Pro Gly Pro Ala Tyr Ser Gly Arg Glu Thr
48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66

                410         420         430         440         450
                |           |           |           |           |
ATA TAC CCC AAT GCA TCC CTG CTG ATC CAG AAC GTC ACC CAG AAT GAC ACA GGA TTC
Ile Tyr Pro Asn Ala Ser Leu Leu Ile Gln Asn Val Thr Gln Asn Asp Thr Gly Phe
67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85

                460         470         480         490         500         510
                |           |           |           |           |           |
TAC ACC CTA CAA GTC ATA AAG TCA GAT CTT GTG AAT GAA GAA GCA ACC GGA CAG TTC
Tyr Thr Leu Gln Val Ile Lys Ser Asp Leu Val Asn Glu Glu Ala Thr Gly Gln Phe
86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104

                520         530         540         550         560         570
                |           |           |           |           |           |
CAT GTA TAC CCG GAG CTG CCC AAG CCC TCC ATC TCC AGC AAC AAC TCC AAC CCC GTG
His Val Tyr Pro Glu Leu Pro Lys Pro Ser Ile Ser Ser Asn Asn Ser Asn Pro Val
105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123
```

```
         580         590         600         610         620
          |           |           |           |           |
GAG GAC AAG GAT GCT GTG GCC TTC ACC TGT GAA CCT GAG GTT CAG AAC ACA ACC TAC
Glu Asp Lys Asp Ala Val Ala Phe Thr Cys Glu Pro Glu Val Gln Asn Thr Thr Tyr
124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141

     630         640         650         660         670         680
      |           |           |           |           |           |
CTG TGG TGG GTA AAT GGT CAG AGC CTC CCG GTC AGT CCC AGG CTG CAG CTG TCC AAT
Leu Trp Trp Val Asn Gly Gln Ser Leu Pro Val Ser Pro Arg Leu Gln Leu Ser Asn
143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161

     690         700         710         720         730         740
      |           |           |           |           |           |
GGC AAC AGG ACC CTC ACT CTA CTC AGC GTC AAA AGG AAC GAT GCA GGA TCG TAT GAA
Gly Asn Arg Thr Leu Thr Leu Leu Ser Val Lys Arg Asn Asp Ala Gly Ser Tyr Glu
162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180

         750         760         770         780         790         800
          |           |           |           |           |           |
TGT GAA ATA CAG AAC CCA GCG AGT GCC AAC CGC AGT GAC CCA GTC ACC CTG AAT GTC
Cys Glu Ile Gln Asn Pro Ala Ser Ala Asn Arg Ser Asp Pro Val Thr Leu Asn Val
181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199

         810         820         830         840         850
          |           |           |           |           |
CTC TAT GGC CCA GAT GGC CCC ACC ATT TCC CCC TCA AAG GCC AAT TAC CGT CCA GGG
Leu Tyr Gly Pro Asp Gly Pro Thr Ile Ser Pro Ser Lys Ala Asn Tyr Arg Pro Gly
200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218

     860         870         880         890         900         910
      |           |           |           |           |           |
GAA AAT CTG AAC CTC TCC TGC CAC GCA GCC TCT AAC CCA CCT GCA CAG TAC TCT TGG
Glu Asn Leu Asn Leu Ser Cys His Ala Ala Ser Asn Pro Pro Ala Gln Tyr Ser Trp
219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237

         920         930         940         950         960         970
          |           |           |           |           |           |
TTT ATC AAT GGG ACG TTC CAG CAA TCC ACA CAA GAG CTC TTT ATC CCC AAC ATC ACT
Phe Ile Asn Gly Thr Phe Gln Gln Ser Thr Gln Glu Leu Phe Ile Pro Asn Ile Thr
238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256

         980         990        1000        1010        1020
          |           |           |           |           |
GTG AAT AAT AGC GGA TCC TAT ATG TGC CAA GCC CAT AAC TCA GCC ACT GGC CTC AAT
Val Asn Asn Ser Gly Ser Tyr Met Cys Gln Ala His Asn Ser Ala Thr Gly Leu Asn
257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275

    1030        1040        1050        1060        1070        1080
     |           |           |           |           |           |
AGG ACC ACA GTC ACG ATG ATC ACA GTC TCT GGA AGT GCT CCT GTC CTC TCA GCT GTG
Arg Thr Thr Val Thr Met Ile Thr Val Ser Gly Ser Ala Pro Val Leu Ser Ala Val
272 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294

    1090        1100        1110        1120        1130        1140
     |           |           |           |           |           |
GCC ACC GTC GGC ATC ACG ATT GGA GTG CTG GCC AGG GTG GCT CTG ATA TAG CAG CCC
Ala Thr Val Gly Ile Thr Ile Gly Val Leu Ala Arg Val Ala Leu Ile ---
295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313
```

1150 1160 1170 1180 1190

TGG TGT ATT TTC GAT ATT TCA GGA AGA CTG GCA GAT TGG ACC AGA CCC TGA ATT CTT

1200 1210 1220 1230 1240 1250

CTA GCT CCT CCA ATC CCA TTT TAT CCC ATG GAA CCA CTA AAA ACA AGG TCT GCT CTG

1260 1270 1280 1290 1300 1310

CTC CTG AAG CCC TAT ATG CTG GAG ATG GAC AAC TCA ATG AAA ATT TAA AGG AAA AAC

1320 1330 1340 1350 1360 1370

CCT CAG GCC TGA GGT GTG TGC CAC TCA GAG ACT TCA CCT AAC TAG AGA CAG GCA AAC

1380 1390 1400 1410 1420

TGC AAA CCA nnC CTC TTT CGC TTG GCA GGA TGA TGG TGT CAT TAG TAT TTC ACA AGA

1430 1440 1450 1460 1470 1480

AGT AGC TTC AGA GGG TAA CTT AAC AGA GTA TCA GAT CTa TCT TGT CAA TCC CAA CGT

1490 1500 1510 1520 1530 1540

TTT ACA TAA AAT AAG CGA TCC TTT AGT GCA CCC AGT GAC TGA CAT TAG CAG CAT CTT

1550 1560 1570 1580 1590

TAA CAC AGC CGT GTG TTC AAG TGT ACA GTG GTC CTT TTC AGA GTT GGn nnT ACT CCA

1600 1610 1620 1630 1640 1650

ACT GAA ATG TTA AGG AAG AAG ATA GAT CCA ATT AAA AAA AAT TAA AAC CAA TTT AAA

1660 1670 1680 1690 1700 1710

AAA AAA AAA GAA CAC AGG AGA TTC CAG TCT ACT TGA GTT AGC ATA ATA CAG AAG TCC

1720 1730 1740 1750 1760

CCT CTA CTT TAA CTT TTA CAA AAA AGT AAC CTG AAC TAA TCT GAT GTT AAC CAA TGT

```
1770        1780        1790        1800        1810        1820
 |           |           |           |           |           |
ATT TAT TTG TCT GGT TCT GTT TCC TTG TTC CAA TTT GAC AAA ACC CAC TGT TCT TGT

1830        1840        1850        1860        1870        1880
 |           |           |           |           |           |
ATT GTA TTG CCC AGG GGG AGC TAT CAC TGT ACT TGT AGA GTG GTG CTG CTT TAA GTT

1890        1900        1910        1920        1930        1940
 |           |           |           |           |           |
CAT AAA TCA CAA ATA AAA GCC AAT TAG CTC TAT AAC TAA AAA AAA AAA AAA AAA AAA

1950        1960
 |           |
AAA AAA AAA AAA AAA AAA AAA AAA
```

A schematic relationship of the transmembrane CEA's, namely TM-1 (CEA-(c)), TM-2 (CEA-(e)), TM-3 (CEA-(f)) and TM-4 (CEA-(g)) is depicted in Fig. 1:

Assuming TM-1 is composed of five sections as depicted in Fig. 1, namely 10, 12, 14, 16 and 18, TM-2 differs from TM-1 in that the 100 amino acid (100 AA) section 14 is deleted and at splice point 20 betweeen sections 12 and 16, surprisingly an extra amino acid, namely Asp occurs.

TM-3 is the same as TM-1 except that section 18 is truncated at splice point 22, i.e., a section of 70 amino acids is deleted and results in a new section made up of subsections 24 + 26. Surprisingly, however, six new amino acids (section 26) occur. Another example of formation of a novel amino acid sequence resulting from a deletion of nucleic acid sequence is for platelet derived growth factor-A.

TM-4 is the same as TM-2 up until the end of subsection 24.

Subsection 24 is contained in section 18 of TM-1 and TM-2, but is not depicted in Fig. 1 for TM-1 and TM-2.

Some CEA epitopes are unique. These are the epitopes which have been useful for distinguishing the various CEA-like antigens immunologically. Peptide epitopes are defined by the linear amino acid sequence of the antigen and/or features resulting from protein folding. The information required for protein folding is encoded in the primary amino acid sequence. Therefore, antigenic differences ultimately result from differences in the primary structure of the different CEA molecules. The differences residing in the CEA protein in the CEA species can thus be determined by determining the primary amino acid sequences. This can be most readily accomplished by cloning and sequencing each of the genes for CEA. To determine which gene products will be most useful for cancer diagnosis, unique probes can be selected for each gene and expression of each gene can be determined in different tumor types by nucleic acid hybridization techniques. The present invention provides a tool with which to identify potential genes coding for different members of the CEA family and to determine the theoretical primary amino acid sequences for them. Using the method of automated peptide synthesis, peptides can then be synthesized corresponding to unique sequences in these antigens. With these peptides, antibodies to these sequences can be produced which, in the intact CEA molecule, might not be recognized by the animal being immunized. Having accomplished this, advantage can then be taken of the differences in these antigens to generate specific immunoassays for the measurement of each antigen.

A wide variety of host/cloning vehicle combinations may be employed in cloning the double-stranded nucleic acid prepared in accordance with this invention. For example, useful cloning vehicles may consist of segments of chromosomal, non-chromosomal and synthetic DNA sequences, such as various known derivatives of SV40 and known bacterial plasmids, e.g., plasmids from E. coli including E1, pCR1, pBR322, pMB89 and their derivatives, wider host range plasmids, e.g., RP4, and phage DNAs, e.g., the numerous derivatives of phage, e.g., NM989, and other DNA phages, e.g., M13 and Filamenteous single-stranded DNA phages and vectors derived from combinations of plasmids and phage DNAs such as plasmids which have been modified to employ phage DNA or other expression control sequences or yeast plasmids such

as the 2 μ plasmid or derivatives thereof. Useful hosts may include bacterial hosts such as strains of E. coli, such as E. coli HB 101, E. coli X1776, E. coli X2282, E. coli MRCI and strains of Pseudomonas, Bacillus subtilis, Bacillus stearothermophilus and other E. coli, bacilli, yeasts and other fungi, animal or plant hosts such as animal (including human) or plant cells in culture or other hosts. Of course, not all host/vector combinations may be equally efficient. The particular selection of host/cloning vehicle combination may be made by those of skill in the art after due consideration of the principles set forth without departing from the scope of this invention.

Furthermore, within each specific cloning vehicle, various sites may be selected for insertion of the nucleic acid according to the present invention. These sites are usually designated by the restriction endonuclease which cuts them. For example, in pBR322 the PstI site is located in the gene for beta-lactamase, between the nucleotide triplets that code for amino acids 181 and 182 of that protein. One of the two HindII endonuclease recognition sites is between the triplets coding for amino acids 101 and 102 and one of the several Taq sites at the triplet coding for amino acid 45 of beta-lactamase in pBR322. In similar fashion, the EcoRI site and the PVUII site in this plasmid lie outside of any coding region, the EcoRI site being located between the genes coding for resistance to tetracycline and ampicillin, respectively. These sites are well recognized by those of skill in the art. It is, of course, to be understood that a cloning vehicle useful in this invention need not have a restriction endonuclease site for insertion of the chosen DNA fragment. Instead, the vehicle could be cut and joined to the fragment by alternative means.

The vector or cloning vehicle and in particular the site chosen therein for attachment of a selected nucleic acid fragment to form a recombinant nucleic acid molecule is determined by a variety of factors, e.g., the number of sites susceptible to a particular restriction enzyme, the size of the protein to be expressed, the susceptibility of the desired protein to proteolytic degradation by host cell enzymes, the contamination of the protein to be expressed by host cell proteins difficult to remove during purification, the expression characteristics, such as the location of start and stop codons relative to the vector sequences, and other factors recognized by those of skill in the art. The choice of a vector and an insertion site for a particular gene is determined by a balance of these factors, not all sections being equally effective for a given case.

Methods of inserting nucleic acid sequences into cloning vehicles to form recombinant nucleic acid molecules include, for example, dA-dT tailing, direct ligation, synthetic linkers, exonuclease and polymerase-linked repair sections followed by ligation, or extension of the nucleic acid strand with an appropriate polymerase and an appropriate single-stranded template followed by ligation.

It should also be understood that the nucleotide sequences or nucleic acid fragments inserted at the selected site of the cloning vehicle may include nucleotides which are not part of the actual structural gene for the desired polypeptide or mature protein or may include only a fragment of the complete structural gene for the desired protein or mature protein.

The cloning vehicle or vector containing the foreign gene is employed to transform an appropriate host so as to permit that host to replicate the foreign gene and to express the protein coded by the foreign gene or portion thereof. The selection of an appropriate host is also controlled by a number of factors recognized by the art. These include, for example, the compatibility with the chosen vector, the toxicity of proteins encoded by the hybrid plasmid, the ease of recovery of the desired protein, the expression characteristics, biosafety and costs. A balance of these factors must be struck with the understanding that not all hosts may be equally effective for expression of a particular recombinant DNA molecule.

The level of production of a protein is governed by two major factors: the number of copies of its gene within the cell and the efficiency with which those gene copies are transcribed and translated. Efficiency of transcription and translation (which together comprise expression) is in turn dependent upon nucleotide sequences, normally situated ahead of the desired coding sequence. These nucleotide sequences or expression control sequences define inter alia, the location at which RNA polymerase interacts to initiate transcription (the promoter sequence) and at which ribosomes bind and interact with the mRNA (the product of transcription) to initiate translation. Not all such expression control sequences function with equal efficiency. It is thus of advantage to separate the specific coding sequences for the desired protein from their adjacent nucleotide sequences and fuse them instead to other known expression control sequences so as to favor higher levels of expression. This having been achieved, the newly engineered nucleic acid, e.g., DNA, fragment may be inserted into a multicopy plasmid or a bacteriophage derivative in order to increase the number of gene copies within the cell and thereby further improve the yield of expressed protein.

Several expression control sequences may be employed as described above. These include the operator, promoter and ribosome binding and interaction sequences (including sequences such as the Shine-Dalgarno sequences) of the lactose operon of E. coli ("the lac system"), the corresponding sequences of the tryptophan synthetase system of E. coli ("the trp system"), the major operator and

promoter regions of phage λ (O_LP_L and $O_RP'_R$), the control region of Filamenteous single-stranded DNA phages, or other sequences which control the expression of genes of prokaryotic or eukaryotic cells and their viruses. Therefore, to improve the production of a particular polypeptide in an appropriate host, the gene coding for that polypeptide may be selected and removed from a recombinant nucleic acid molecule containing it and reinserted into a recombinant nucleic acid molecule closer or in a more appropriate relationship to its former expression control sequence or under the control of one of the above described expression control sequences. Such methods are known in the art.

As used herein "relationship" may encompass many factors, e.g., the distance separating the expression enhancing and promoting regions of the recombinant nucleic acid molecule and the inserted nucleic acid sequence, the transcription and translation characteristics of the inserted nucleic acid sequence or other sequences in the vector itself, the particular nucleotide sequence of the inserted nucleic acid sequence and other sequences of the vector and the particular characteristics of the expression enhancing and promoting regions of the vector.

Further increases in the cellular yield of the desired products depend upon an increase in the number of genes that can be utilized in the cell. This is achieved, for illustration purposes, by insertion of recombinant nucleic acid molecules engineered into the temperate bacteriophageλ (NM989), most simply by digestion of the plasmid with a restriction enzyme, to give a linear molecule which is then mixed with a restricted phage λ cloning vehicle (e.g., of the type described by N. E. Murray et al, "Lambdoid Phages That Simplify the Recovery of In Vitro Recombinants", Molec. Gen. Genet., 150, pp. 53-61 (1977) and N.E. Murray et al, "Molecular Cloning of the DNA Ligase Gene From Bacteriophage T4", J. Mol. Biol., 132, pp. 493-505 (1979)) and the recombinant DNA molecule recircularized by incubation with DNA ligase. The desired recombinant phage is then selected as before and used to lysogenize a host strain of E. coli.

Particularly useful λ cloning vehicles contain a temperature-sensitive mutation in the repression gene cI and suppressible mutations in gene S, the product of which is necessary for lysis of the host cell, and gene E, the product of which is major capsid protein of the virus. With this system, the lysogenic cells are grown at 32° C and then heated to 45° C to induce excision of the prophage. Prolonged growth at 37° C leads to high levels of production of the protein, which is retained within the cells, since these are not lysed by phage gene products in the normal way, and since the phage gene insert is not encapsulated it remains available for further transcription. Artificial lysis of the cells then releases the desired product in high yield.

In addition, it should be understood that the yield of polypeptides prepared in accordance with this invention may also be improved by substituting different codons for some or all of the codons of the present DNA sequences, these substituted codons coding for amino acids identical to those coded for by the codons replaced.

Finally, the activity of the polypeptides produced by the recombinant nucleic acid molecules of this invention may be improved by fragmenting, modifying or derivatizing the nucleic acid sequences or polypeptides of this invention by well-known means, without departing from the scope of this invention.

The polypeptides of the present invention include the following:

(1) the polypeptides expressed by the above described cells,

(2) polypeptides prepared by synthetic means,

(3) fragments of polypeptides (1) or (2) above, such fragments produced by synthesis of amino acids or by digestion or cleavage.

Regarding the synthetic peptides according to the invention, chemical synthesis of peptides is described in the following publications: S.B.H. Kent, Biomedical Polymers, eds. Goldberg, E.P. and Nakajima, A. (Academic Press, New York), 213-242, (1980); A.R. Mitchell, S.B.H. Kent, M. Engelhard and R.B. Merrifield, J. Org. Chem., 43, 2845-2852, (1978); J.P. Tam, T.-W. Wong, M. Rieman, F.-S. Tjoeng and R.B. Merrifield, Tet. Letters, 4033-4036, (1979); S. Mojsov, A.R. Mitchell and R.B. Merrifield, J. Org. Chem., 45, 555-560, (1980); J.P. Tam, R.D. DiMarchi and R.B. Merrifield, Tet. Letters, 2851-2854, (1981); and S.B.H. Kent, M. Riemen, M. Le Doux and R.B. Merrifield, Proceedings of the IV International Symposium on Methods of Protein Sequence Analysis, (Brookhaven Press, Brookhaven, NY), in press, 1981.

In the Merrifield solid phase procedure, the appropriate sequence of L-amino acids is built up from the carboxyl terminal amino acid to the amino terminal amino acid. Starting with the appropriate carboxyl terminal amino acid attached to a polystyrene (or other appropriate) resin via chemical linkage to a chloromethyl group, benzhydrylamine group, or other reactive group of the resin, amino acids are added one by one using the following procedure. The peptide-resin is:

(a) washed with methylene chloride;

(b) neutralized by making for 10 minutes at room temperature with 5% (v/v) diisopropylethylamine (or other hindered base) in methylene chloride;

(c) washed with methylene chloride;

(d) an amount of amino acid equal to six times the molar amount of the growing peptide chain is activated by combining it with one-half as many moles of a carbodiimide (e.g., dicyclohexylcarbodiimide, or diisopropylcarbodiimide) for ten minutes at 0°C, to form the symmetric anhydride of the amino acid. The amino acid used should be provided originally as the N-alpha-tert.-butyloxycarbonyl derivative, with side chains protected with benzyl esters (e.g., aspartic or glutamic acids), benzyl ethers (e.g., serine, threonine, cysteine or tyrosine), benzyloxycarbonyl groups (e.g., lysine) or other protecting groups commonly used in peptide synthesis;

(e) the activated amino acid is reacted with the peptide-resin for two hours at room temperature, resulting in addition of the new amino acid to the end of the growing peptide chain;

(f) the peptide-resin is washed with methylene chloride;

(g) the N-alpha-(tert.-butyloxycarbonyl) group is removed from the most recently added amino acid by reacting with 30 to 65%, preferably 50% (v/v) trifluoroacetic acid in methylene chloride for 10 to 30 minutes at room temperature;

(h) the peptide-resin is washed with methylene chloride;

(i) steps (a) through (h) are repeated until the required peptide sequence has been constructed.

The peptide is then removed from the resin and simultaneously the side-chain protecting groups are removed, by reaction with anhydrous hydrofluoric acid containing 10% v/v of anisole or other suitable (aromatic) scavenger. Subsequently, the peptide can be purified by gel filtration, ion exchange, high pressure liquid chromatography, or other suitable means.

In some cases, chemical synthesis can be carried out without the solid phase resin, in which case the synthetic reactions are performed entirely in solution. The reactions are similar and well known in the art, and the final product is essentially identical.

Digestion of the polypeptide can be accomplished by using proteolytic enzymes, especially those enzymes whose substrate specificity results in cleavage of the polypeptide at sites immediately adjacent to the desired sequence of amino acids.

Cleavage of the polypeptide can be accomplished by chemical means. Particular bonds between amino acids can be cleaved by reaction with specific reagents. Examples include the following: bonds involving methionine are cleaved by cyanogen bromide; asparaginyl-glycine bonds are cleaved by hydroxylamine.

The present invention has the following advantages:

(1) The nucleic acids coding for TM-1, TM-2 and TM-3 can be used as probes to isolate other members of the CEA gene family.

(2) The nucleic acids coding for TM-1, TM-2 and TM-3 can be used to derive oligonucleotide probes to determine the expression of TM-1, TM-2, TM-3 and other CEA genes in various tumor types.

(3) TM-1, TM-2, TM-3 and TM-4 nucleotide sequences can be used to predict the primary amino acid sequence of the protein for production of synthetic peptides.

(4) Synthetic peptides derived from the above sequences can be used to produce sequence-specific antibodies.

(5) Immunoassays for each member of the CEA antigen family can be produced with these sequence-specific antibodies and synthetic peptides.

(6) The aforementioned immunoassays can be used as diagnostics for different types of cancer if it is determined that different members of the CEA family are clinically useful markers for different types of cancer.

Peptides according to the present invention can be labelled by conventional means using radioactive moieties (e.g., ^{125}I), enzymes, dyes and fluorescent compounds, just to name a few.

Several possible configurations for immunoassays according to the present invention can be used. The readout systems capable of being employed in these assays are numerous and non-limiting examples of such systems include fluorescent and colorimetric enzyme systems, radioisotopic labelling and detection and chemiluminescent systems. Two examples of immunoassay methods are as follows:

(1) An enzyme linked immunoassay (ELISA) using an antibody preparation according to the present invention (including Fab or F(ab)' fragments derived therefrom) to a solid phase (such as a microtiter plate or latex beads) is attached a purified antibody of a specificity other than that which is conjugated to the enzyme. This solid phase antibody is contacted with the sample containing CEA antigen family members. After washing, the solid phase antibody-antigen complex is contacted with the conjugated anti-peptide antibody (or conjugated fragment). After washing away unbound conjugate, color or fluorescence is developed by adding a chromogenic or fluorogenic substrate for the enzyme. The amount of color or fluorescence developed is proportional to the amount of antigen in the sample.

(2) A competitive fluorometric immunoassay using fluorescently labelled peptide or synthetic peptides of the sequences for TM-2, TM-2, TM-3 and TM-4. In this example, the purified peptide expressed by cells or synthetic peptides thereof are fluorescently labelled. To a solid phase is attached a purified antibody. This solid phase is then contacted with sample containing CEA antigen family members to which has been added fluorescent peptide probe. After binding, excess probe is washed away the amount of bound probe is quantitated. The amount of bound fluorescent probe will be inversely proportional to the amount of antigen in the sample.

In the nucleic acid hybridization method according to the present invention, the nucleic acid probe is conjugated with a label, for example, an enzyme, a fluorophore, a radioisotope, a chemiluminescent compound, etc. In the most general case, the probe would be contacted with the sample and the presence of any hybridizable nucleic acid sequence would be detected by developing in the presence of a chromogenic enzyme substrate, detection of the fluorophore by epifluorescence, by autoradiography of the radioisotopically labelled probe or by chemiluminescence. The detection of hybridizable RNA sequences can be accomplished by (1) a dot blot methodology or (2) an in situ hybridization methodology. Methods for these last two techniques are described by D. Gillespie and J. Bresser, "mRNA Immobilization in NaI: Quick Blots", Biotechniques, 184-192, November/December 1983 and J. Lawrence and R. Singer, "Intracellular Localization of Messenger RNAs for Cytosketal Proteins", Cell, 45, 407-415, May 9, 1986, respectively. The readout systems can be the same as described above, e.g., enzyme labelling, radiolabelling, etc.

As stated above, the invention also relates to the use in medicine of the aforementioned complex of the invention.

The invention further provides a pharmaceutical composition containing as an active ingredient a complex of the invention in the form of a sterile and/or physiologically isotonic aqueous solution.

For parenteral administration, solutions and emulsions containing as an active ingredient the complex of the invention should be sterile and, if appropriate, blood-isotonic.

It is envisaged that the active complex will be administered perorally, parenterally (for example, intramuscularly, intraperitoneally, or intravenously), rectally or locally.

Example 1: Preparation of cDNA in pcE22 which codes for TM2-CEA [CEA-(e)]

Example 1a: RNA Preparation

Messenger RNA was prepared by the proteinase K extraction method of J. Favolaro, R. Treisman and R. Kamen, Methods in Enzymology, 65, 718, Academic Press, Inc. (1980), followed by oligo dT cellulose chromatography to yield poly A+ RNA (3′-polyadenylated eukaryotic RNA containing most mRNA sequences that can be translated into polypeptides). To obtain approximately 100 µg of poly A+ RNA, approximately 3 x 10^8 cells of transfectant 23.411 (ATCC No. CRL 9731, deposited with the ATCC on June 1, 1988), that expresses TM-1, TM-2, TM-3 and TM-4, Kamarck et al, Proc. Natl. Acad. Sci., USA, 84, 5350-5354, August 1987, were harvested from roller bottles after late logarithmic growth. Cells were lysed by homogenization in an ice-cold solution of 140 mM NaCl, 1.5 mM $MgCl_2$, 10 mM Tris-HCl, pH 8.0, 0.5% NP40, 4 mM dithiothreitol and 20 units of placental ribonuclease inhibitor/ml. sodium deoxycholate was then added to 0.2%. Cytoplasm and nuclei were separated by centrifugation of the homogenate at 12,000xg for 20 minutes. The cytoplasmic fraction was mixed with an equal volume of 0.2 M Tris-HCl, pH 7.8, 25 mM EDTA, 0.3 M NaCl, 2% sodium dodecyl sulfate and 400 µg/ml of proteinase K, incubated for 1 hour at 37° C, then extracted once with an equal volume of phenol/chloroform (1:1/v:v) solution. Nucleic acids were obrtained by ethanol precipitation of the separated aqueous phase. Total RNA was enriched by passage in 0.5 M NaCl, 10 mM Tris-HCl, pH 7.8, 0.1% sarcosyl through an oligo dT(12-18) cellulose column. After washing, bound RNA was eluted in the same solution without sodium chloride.

Example 1b: Reverse Transcription of mRNA

Ten micrograms of poly A+ RNA were primed for reverse transcription with oligo dT(12-18) and pdN_6 primers. One hundred microliter reaction was performed for 4 hours at 42° C with 200 units AMV reverse transcriptase (Life Science, Inc. St. Petersburg, Florida, U.S.A.). The RNA component of the cDNA/mRNA hybrids was replaced with the second complementary strand by treatment with RNase H, E. coli DNA

polymerase I and E. coli DNA ligase at 12° C and 22° C for 1.5 hours each. Molecular ends were polished by treatment with T4 DNA polymerase. cDNA was phenol/chloroform extracted and purified over a "SEPHADEX G-50" spun column prepared in 10 mM Tris-HCl, pH 7.8, 1 mM EDTA (TE).

Example 1c: Cloning of pcE23 (plasmid cDNA E22)

Synthetic DNA linkers

```
5' pCCCGGG     3'
3'  GGGCCCTTAA 5'
```

were attached to the ends of cDNA by blunt end ligation with excess T4 DNA ligase. Excess linkers were removed by chromatography through "SEPHADEX G-50" (medium) in TE, and by fractionation on 0.8% low melting agarose gel. Based on Northern blot analysis of poly A+ RNA of the 23.411 cell line, the size of the CEA-related mRNA was estimated at 3.6 kb. Therefore, cDNA fragments between 2 and 4 kb were recovered from gel slices and fragments were ethanol precipitated. After resuspension of cDNA in TE, EcoRI-cleaved lambada gt10 arms were added to cDNA at an estimated molar ration of 1:1. Ligation proceeded at 7° C for 2 days in the presence of T4 DNA ligase. Aliquots of the ligation reaction were added to commercially-obtained packaging mix (Stratagene, San Diego, California, U.S.A.). Five million phage particles were obtained ofter in vitro packaging and infection of E. coli host NM514.

Example 1d: Screening of Recombinant Library

Five hundred thousand packaged lambda particles were plated on lawns of E. coli NM514 and replicate patterns were lifted onto nitrocellulose sheets as described by W.D. Benton and R.W. Davis, Science 196, 180-182, (1977). Positive phage were selected by hybridization with ^{32}P-labeled LV7 cDNA insert probe that contained a domain repeated amoung various CEA family members. By multiple rounds of screening. Phage from individual plaques were amplified and titered, and these were used to prepare small quantities of recombinant phage DNA.

Example 1e: DNA Manipulation

Phage DNA was prepared according to T. Maniatis, E. Fritsch and J. Sambrook, Molecular Cloning, A Laboratory Manual, Cold Spring Habor, (1982). DNA segments were isolated from low melting agarose gels and inserted for subcloning into Bluescript plasmid vectors (Stratagene, San Diego, California, U.S.A.). DNA sequencing was performed by the dideoxy termination method of F. Sanger, S. Nicklen and A. Coulson, Proc. Natl. Acad. Sci., U.S.A., 74, 5463-5467, (1977). The nucleic acid and translated sequence for cDNA in pcE22 is given hereinabove (TM-2 (CEA-(e)).

Example 2: Preparation of cDNA in pcHT-6 which Partially Codes for TM3-CEA [CEA-(f)]

Example 2a: RNA Preparation

Messenger RNA was prepared by the proteinase K extraction method of J. Favolaro, R. Treisman and R. Kamen, Methods in Enzymology, 65 718, Academic Press, Inc. (1980), followed by oligo dT cellulose chromatography to yield poly A+ RNA (3'-polyadenylated eukaryotic RNA containing most mRNA sequences that can be translated into polypeptides). To obtain approximately 100 ug of poly A+ RNA, approximately 3 x 10^8 cells of HT-29 tumor cells (ATCC HTB38) were harvested form roller bottles after late logarithmic growth. Cells were lysed by homogenization in an ice-cold solution of 140 mM NaCl, 1.5 mM $MgCl_2$, 10 mM Tris-HCl, pH 8.0, 0.5% NP40, 4 mM dithiothreitol and 20 units of placental ribonuclease inhibitor/ml. Sodium deoxycholate was then added to 0.2%. Cytoplasm and nuclei were separated by

centrifugation of the homogenate at 12,000xg for 20 minutes. The cytoplasmic fraction was mixed with an equal volume of 0.2 M Tris-Hcl, pH 7.8, 25 mM EDTA, 0.3 M NaCl, 2% sodium dodecyl sulfate and 400 µg/ml of proteinase K, incubated for 1 hour at 37° C, then extracted once with an equal volume of phenol/chloroform (1:1/v:v) solution. Nucleic acids were obrtained by ethanol precipitation of the separated aqueous phase. Total RNA was enriched by passage in 0.5 M NaCl, 10 mM Tris-HCl, pH 7.8, 0.1% sarcosyl through an oligo dT(12-18) cellulose column. After washing, bound RNA was eluted in the same solution without sodium chloride.

Example 2b: Reverse Transcription of mRNA

Ten micrograms of HT-29 poly A+ RNA were primed for reverse transcription with oligo dT(12-18) and pdN_6 primers. One hundred microliter reaction was performed for 4 hours at 42° C with 200 units AMV reverse transcriptase (Life Science, Inc. St. Petersburg, Florida, U.S.A.). The RNA component of the cDNA/mRNA hybrids was replaced with the second complementary strand by treatment with RNase H, E. coli DNA polymerase I and E. coli DNA ligase at 12° C and 22° C for 1.5 hours each. Molecular ends were polished by treatment with T4 DNA polymerase. cDNA was phenol/chloroform extracted and purified over a "SEPHADEX G-50" spun column prepared in 10 mM Tris-HCl, pH 7.8, 1 mM EDTA (TE).

Example 2c: Cloning of pcHT-6 (plasmid cDNA HT-6)

Synthetic DNA linkers

```
5' pCCCGGG    3'
3'  GGGCCCTTAA 5'
```

were attached to the ends of cDNA by blunt end ligation with excess T4 DNA ligase. Excess linkers were removed by chromatography through "SEPHADEX G-50" (medium) in TE, and by fractionation on 0.8% low melting agarose gel. Based on Northern blot analysis of poly A+ RNA of the HT-29 cell line, the size of the CEA-related mRNA was estimated at 2.2 kb. Therefore, cDNA fragments between 2 and 3 kb were recovered from gel slices and fragments were ethanol precipitated. After resuspension of cDNA in TE, EcoRI-cleaved lambada gt10 arms were added to cDNA at an estimated molar ration of 1:1. Ligation proceeded at 7° C for 2 days in the presence of T4 DNA ligase. Aliquots of the ligation reaction were added to commercially-obtained packaging mix (Stratagene, San Diego, California, U.S.A.). Five million phage particles were obtained ofter in vitro packaging and infection of E. coli host NM514.

Example 2d: Screening of Recombinant Library

Five hundred thousand packaged lambda particles were plated on lawns of E. coli NM514 and replicate patterns were lifted onto nitrocellulose sheets as described by W.D. Benton and R.W. Davis, Science, 196, 180-182, (1977). Positive phage were selected by hybridization with ^{32}P-labeled LV7 cDNA insert probe that contained a domain repeated amoung various CEA family members. By multiple rounds of screening. Phage from individual plaques were amplified and titered, and these were used to prepare small quantities of recombinant phage DNA.

Example 2e: DNA Manipulation

Phage DNA was prepared according to T. Maniatis, E. Fritsch and J. Sambrook, Molecular Cloning, A Laboratory Manual, Cold Spring Habor, (1982). DNA segments were isolated from low melting agarose gels and inserted for subcloning into Bluescript plasmid vectors (Stratagene, San Diego, California, U.S.A.). DNA sequencing was performed by the dideoxy termination method of F. Sanger, S. Nicklen and A. Coulson, Proc. Natl. Acad. Sci., U.S.A., 74, 5463-5467, (1977). The nucleic acid and translated sequence for cDNA in HT-6 not complete at the 5' end of its coding region, but the nucleotide sequenece and restriction map of

the HT-6 insert indicates that it is related to nucleic acid sequences of cDNA clones coding for CEA-(c) and CEA-(e). The nucleotide sequence of HT-6 insert differs from these clones at only nucleotide position 1463 to 1515 and 1676 to 2429 of the CEA-(c) cDNA. It is inferred from this result that the pcHT-6 insert is a partial coding sequence for CEA-(f) and the presumed nucleic acid and translated sequence of CEA-(f) is given hereinabove (TM-3 (CEA-(f)).

Example 3: Preparation of cDNA which codes for TM4-CEA [CEA-(g)]

Example 3a: RNA Preparation

Messenger RNA is prepared by the proteinase K extraction method of J. Favolaro, R. Treisman and R. Kamen, Methos in Enzymology, 65, 718, Academic Press, Inc. (1980), followed by oligo dT cellulose chromatography to yield poly A+ RNA (3'-polyadenylated eukaryotic RNA containing most mRNA sequences that can be translated into polypeptides). To obtain approximately 100 ug of poly A+ RNA, approximately 3 x 10^8 cells of transfectant 23.411 or tumor cell line HT-29 (ATCC HTB 38) are harvested from roller bottles after late logarithmic growth. Cells are lysed by homogenization in an ice-cold solution of 140 mM NaCl, 1.5 mM $MgCl_2$, 10 mM Tris-HCl, pH 8.0, 0.5% NP40, 4 mM dithiothreitol and 20 units of placental ribonuclease inhibitor/ml. Sodium deoxycholate was then added to 0.2%. Cytoplasm and nuclei are separated by centrifugation of the homogenate at 12,000xg for 20 minutes. The cytoplasmic fraction is mixed with an equal volume of 0.2 M Tris-Hcl, pH 7.8, 25 mM EDTA, 0.3 M NaCl, 2% sodium dodecyl sulfate and 400 µg/ml of proteinase K, incubated for 1 hour at 37° C, then extracted once with an equal volume of phenol/cholorform (1:1/v:v) solution. Nucleic acids are obtained by ethanol precipitation of the separated aqueous phase. Total RNA is enriched by passage in 0.5 M NaCl, 10 mM Tris-HCl, pH 7.8, 0.1% sarcosyl through an oligo dT(12-18) cellulose column. After washing, bound RNA is eluted in the same solution without sodium chloride.

Example 3b: Reverse Transcription of mRNA

Ten micrograms of 23.411 or HT 29 poly A+ RNA are primed for reverse transcription with oligo dT-(12-18) and pdN_6 primers. One hundred microliter reaction was performed for 4 hours at 42° C with 200 units AMV reverse transcriptase (Life Science, Inc. St. Petersburg, Florida, U.S.A.). The RNA component of the cDNA/mRNA hybrids is replaced with the second complementary strand by treatment with RNase H, E. coli DNA polymerase I and E. coli DNA ligase at 12° C and 22° C for 1.5 hours each. Molecular ends are polished by treatment with T4 DNA polymerase. cDNA is phenol/chloroform extracted and purified over a "SEPHADEX G-50" spun column prepared in 10 mM Tris-HCl, pH 7.8, 1 mM EDTA (TE).

Example 3c: Cloning of cDNA for CEA-(g)

Synthetic DNA linkers

```
5' pCCCGGG    3'
3'  GGGCCCTTAA 5'
```

are attached to the ends of cDNA by blunt end ligation which excess T4 DNA ligase. Excess linkers are removed by chromatography through "SEPHADEX G-50" (medium) in TE, and by fractionation on 0.8% low melting agarose gel. Based on Northern blot analysis of poly A+ RNA of the 23.411 and HT-29 cell lines, the size of the CEA-related mRNA is estimated at 1.7 kb. Therefore, cDNA fragments between 1 and 2 kb are recovered from gel slices and fragments are ethanol precipitated. After resuspension of cDNA in TE, EcoRI-cleaved lambda gt10 arms are added to cDNA at an estimated molar ratio of 1:1. Ligation proceeds at 7° C for 2 days in the presence of T4 DNA ligase. Aliquots of the ligation reaction are added to commercially-obtained packaging mix (Stratagene, San Diego, California, U.S.A.). Phage particles are obtained after in vitro packaging and infection of E. coli host NM514.

Example 3d: Screening of Recombinant Library

Five hundred thousand to one million packaged lambda particles are plated on lawns of E. coli NM514 and replicate patterns are lifted onto nitrocellulose sheets as described by W.D. Benton and R.W. Davis, Science, 196, 180-182, (1977). Positive phage are selected by hybridization with ^{32}P-labeled LV7 cDNA insert probe that contained a domain repeated amoung various CEA family members. By this selection method, positive phage are obtained after multiple rounds of screening. Phage from individual plaques are amplified and titered, and these are used to prepare small quantities of recombinant phage DNA.

Example 3e: DNA Manipulation

Phage DNA is prepared according to T. Maniatis, E. Fritsch and J. Sambrook, Molecular Cloning, A Laboratory Manual, Cold Spring Harbor, (1982). DNA segments are isolated from low melting agarose gels and inserted for subcloning into Bluescript plasmid vectors (Stratagene, San Diego, California, U.S.A.). DNA sequencing is performed by the dideoxy termination method of F. Sanger, S. Nicklen and A. Coulson, Proc. Natl. Acad. Sci., U.S.A., 74, 5463-5467, (1977). The nucleotide and translated sequence for a cDNA coding for CEA-(g) is given hereinabove (TM-4 (CEA-(g)).

Example 4: Screening of a KG-1 cDNA Library with ^{32}P-labelled CEA Probe, LV7 (CEA-(A))

A segment of cDNA coding for a portion of carcinoembryonic antigen [LV7 or CEA-(a)] was radiolabelled by random priming and used to detect homologous sequences on filter replicas of a commercial cDNA library prepared from KG-1 cells in bacteriophage vector λ gt11 (Clontech Laboratories, Inc., Palo Alto, CA., U.S.A.). Hybridizations were performed at 68° C in 2xSSSPE (1xSSPE - 0.15 M NaCl, 0.01 M sodium phosphate and 1 mM EDTA, pH 7), 5x Denhardt's solution and 100 µg of denatured salmon sperm DNA per ml, and post-hybridization washes were in 0.2xSSC, 0.25% sodium dodecyl sulfate.

Positive phage were picked, rescreened to homogeneity, and amplified for production of DNA. cDNA inserts were excised from phage DNA with EcoRI endonuclease and subcloned into the EcoRI site of the plasmid vector pBluescript KS. DNA sequencing on double-stranded DNA was by the method of Sanger et al, supra. The sequences of two different inserts from the KG-1 cDNA library are shown below:

pcKGCEA1:

```
    1  acagcacagctgacagccgtactcaggaagcttctggatcctaggcttatctccacagag   60

   61  gagaacacacaagcagcagagaccatggggcccctctcagcccctccctgcacacacctc  120
                                  MetGlyProLeuSerAlaProProCysThrHisLeu

  121  atcacttggaaggggtcctgctcacagcatcacttttaaacttctggaatccgcccaca   180
       IleThrTrpLysGlyValLeuLeuThrAlaSerLeuLeuAsnPheTrpAsnProProThr

  181  actgcccaagtcacgattgaagcccagccacccaaagtttctgaggggaaggatgttctt  240
       ThrAlaGlnValThrIleGluAlaGlnProProLysValSerGluGlyLysAspValLeu

  241  ctacttgtccacaatttgccccagaatcttgctggctacatttggtacaaagggcaaatg  300
       LeuLeuValHisAsnLeuProGlnAsnLeuAlaGlyTyrIleTrpTyrLysGlyGlnMet

  301  acatacgtctaccattacattacatcatatgtagtagacggtcaaagaattatatatggg  360
       ThrTyrValTyrHisTyrIleThrSerTyrValValAspGlyGlnArgIleIleTyrGly

  361  cctgcatacagtggaagagaaagagtatattccaatgcatccctgctgatccagaatgtc  420
       ProAlaTyrSerGlyArgGluArgValTyrSerAsnAlaSerLeuLeuIleGlnAsnVal

  421  acgcaggaggatgcaggatcctacaccttacacatcataaagcgacgcgatgggactgga  480
       ThrGlnGluAspAlaGlySerTyrThrLeuHisIleIleLysArgArgAspGlyThrGly

  481  ggagtaactggacatttcaccttcaccttacacctggagactcccaagccctccatctcc  540
       GlyValThrGlyHisPheThrPheThrLeuHisLeuGluThrProLysProSerIleSer

  541  agcagcaacttaaatcccagggaggccatggaggctgtgatcttaacctgtgatcctgcg  600
       SerSerAsnLeuAsnProArgGluAlaMetGluAlaValIleLeuThrCysAspProAla

  601  actccagccgcaagctaccagtggtggatgaatggtcagagcctccctatgactcacagg  660
       ThrProAlaAlaSerTyrGlnTrpTrpMetAsnGlyGlnSerLeuProMetThrHisArg

  661  ttgcagctgtccaaaaccaacaggaccctctttatatttggtgtcacaaagtatattgca  720
       LeuGlnLeuSerLysThrAsnArgThrLeuPheIlePheGlyValThrLysTyrIleAla

  721  ggaccctatgaatgtgaaatacggaacccagtgagtgccagccgcagtgacccagtcacc  780
       GlyProTyrGluCysGluIleArgAsnProValSerAlaSerArgSerAspProValThr

  781  ctgaatctcctcccaaagctgtccaagccctacatcacaatcaacaacttaaaccccaga  840
       LeuAsnLeuLeuProLysLeuSerLysProTyrIleThrIleAsnAsnLeuAsnProArg

  841  gagaataaggatgtcttaaccttcacctgtgaacctaagagtgagaactacacctacatt  900
       GluAsnLysAspValLeuThrPheThrCysGluProLysSerGluAsnTyrThrTyrIle

  901  tggtggctaaatggtcagagcctccctgtcagtcccagggtaaagcgacccattgaaaac  960
       TrpTrpLeuAsnGlyGlnSerLeuProValSerProArgValLysArgProIleGluAsn

  961  aggatcctcattctacccaatgtcacgagaaatgaaacaggaccttatcaatgtgaaata 1020
       ArgIleLeuIleLeuProAsnValThrArgAsnGluThrGlyProTyrGlnCysGluIle

 1021  cgggaccgatatggtggcatccgcagtgacccagtcaccctgaatgtcctctatggtcca 1080
       ArgAspArgTyrGlyGlyIleArgSerAspProValThrLeuAsnValLeuTyrGlyPro
```

```
1081  gacctccccagcatttacccttcattcacctattaccgttcaggagaaaacctctacttt  1140
      AspLeuProSerIleTyrProSerPheThrTyrTyrArgSerGlyGluAsnLeuTyrPhe

1141  tcctgcttcggtgagtctaacccacgggcacaatattcttggacaattaatgggaagttt  1200
      SerCysPheGlyGluSerAsnProArgAlaGlnTyrSerTrpThrIleAsnGlyLysPhe

1201  cagctatcaggacaaaagctctctatcccccaaataactacaaagcatagtgggctctat  1260
      GlnLeuSerGlyGlnLysLeuSerIleProGlnIleThrThrLysHisSerGlyLeuTyr

1261  gcttgctctgttcgtaactcagccactggcaaggaaagctccaaatccatcacagtcaaa  1320
      AlaCysSerValArgAsnSerAlaThrGlyLysGluSerSerLysSerIleThrValLys

1321  gtctctgactggatattaccctgaattctactagttcctccaattccatttttctcccatg  1380
      ValSerAspTrpIleLeuProEnd

1381  gaatcacgaagagcaagacccactctgttccagaagccctataatctggaggtggacaac  1440
1441  tcgatgtaaatttcatgggaaaaccctttgtacctgacatgtgagccactcagaactcacc  1500
1501  aaaatgttcgacaccataacaacagctactcaaactgtaaaccaggataagaagttgatg  1560
1561  acttcacactgtggacagtttttcaaagatgtcataacaagactccccatcatgacaagg  1620
1621  ctccaccctctactgtctgctcatgcctgcctctttcacttggcaggataatgcagtcat  1680
1681  tagaatttcacatgtagtagcttctgagggtaacaacagagtgtcagatatgtcatctca  1740
1741  acctcaaacttttacgtaacatctcagggaaatgtggctctctccatcttgcatacaggg  1800
1801  ctcccaatagaaatgaacacagagatattgcctgtgtgtttgcagagaagatggtttcta  1860
1861  taaagagtaggaaagctgaaattatagtagagtctcctttaaatgcacattgtgtggatg  1920
1921  gctctcaccatttcctaagagatacagtgtaaaaacgtgacagtaatactgattctagca  1980
1981  gaataaacatgtaccacatttgcaaaaaa                                  2010
```

pcKGCEA2:

```
1     gggtggatcctaggctcatctccataggggagaacacacatacagcagagaccatggga    59
                                                          MetGly

60    ccctctcagcccctccctgcactcagcacatcacctggaaggggctcctgctcacagca   119
      ProLeuSerAlaProProCysThrGlnHisIleThrTrpLysGlyLeuLeuLeuThrAla

120   tcacttttaaacttctggaacctgcccaccactgcccaagtaataattgaagcccagcca   179
      SerLeuLeuAsnPheTrpAsnLeuProThrThrAlaGlnValIleIleGluAlaGlnPro

180   cccaaagtttctgaggggaaggatgttcttctacttgtccacaatttgccccagaatctt   239
      ProLysValSerGluGlyLysAspValLeuLeuLeuValHisAsnLeuProGlnAsnLeu

240   actggctacatctggtacaaagggcaaatgacggacctctaccattacattacatcatat   299
      ThrGlyTyrIleTrpTyrLysGlyGlnMetThrAspLeuTyrHisTyrIleThrSerTyr

300   gtagtagacggtcaaattatatatgggcctgcctacagtggacgagaaacagtatattcc   359
      ValValAspGlyGlnIleIleTyrGlyProAlaTyrSerGlyArgGluThrValTyrSer

360   aatgcatccctgctgatccagaatgtcacacaggaggatgcaggatcctacaccttacac   419
      AsnAlaSerLeuLeuIleGlnAsnValThrGlnGluAspAlaGlySerTyrThrLeuHis

420   atcataaagcgaggcgatgggactggaggagtaactggatatttcactgtcaccttatac   479
      IleIleLysArgGlyAspGlyThrGlyGlyValThrGlyTyrPheThrValThrLeuTyr

480   tcggagactcccaagcgctccatctccagcagcaacttaaaccccagggaggtcatggag   539
      SerGluThrProLysArgSerIleSerSerSerAsnLeuAsnProArgGluValMetGlu
```

```
540  gctgtgcgcttaatctgtgatcctgagactccggatgcaagctacctgtggttgctgaat   599
     AlaValArgLeuIleCysAspProGluThrProAspAlaSerTyrLeuTrpLeuLeuAsn

600  ggtcagaacctccctatgactcacaggttgcagctgtccaaaaccaacaggaccctctat   659
     GlyGlnAsnLeuProMetThrHisArgLeuGlnLeuSerLysThrAsnArgThrLeuTyr

660  ctatttggtgtcacaaagtatattgcagggccctatgaatgtgaaatacggaggggagtg   719
     LeuPheGlyValThrLysTyrIleAlaGlyProTyrGluCysGluIleArgArgGlyVal

720  agtgccagccgcagtgacccagtcaccctgaatctcctcccgaagctgcccatgccttac   779
     SerAlaSerArgSerAspProValThrLeuAsnLeuLeuProLysLeuProMetProTyr

780  atcaccatcaacaacttaaaccccagggagaagaaggatgtgttagccttcacctgtgaa   839
     IleThrIleAsnAsnLeuAsnProArgGluLysLysAspValLeuAlaPheThrCysGlu

840  cctaagagtcggaactacacctacatttggtggctaaatggtcagagcctcccggtcagt   899
     ProLysSerArgAsnTyrThrTyrIleTrpTrpLeuAsnGlyGlnSerLeuProValSer

900  ccgagggtaaagcgacccattgaaaacaggatactcattctacccagtgtcacgagaaat   959
     ProArgValLysArgProIleGluAsnArgIleLeuIleLeuProSerValThrArgAsn

960  gaaacaggaccctatcaatgtgaaatacgggaccgatatggtggcatccgcagtaaccca   1019
     GluThrGlyProTyrGlnCysGluIleArgAspArgTyrGlyGlyIleArgSerAsnPro

1020 gtcaccctgaatgtcctctatggtccagacctccccagaatttacccttacttcacctat   1079
     ValThrLeuAsnValLeuTyrGlyProAspLeuProArgIleTyrProTyrPheThrTyr

1080 taccgttcaggagaaaacctcgacttgtcctgctttgcggactctaacccaccggcagag   1139
     TyrArgSerGlyGluAsnLeuAspLeuSerCysPheAlaAspSerAsnProProAlaGlu

1140 tatttttggacaattaatgggaagtttcagctatcaggacaaaagctctttatcccccaa   1199
     TyrPheTrpThrIleAsnGlyLysPheGlnLeuSerGlyGlnLysLeuPheIleProGln

1200 attactacaaatcatagcgggctctatgcttgctctgttcgtaactcagccactggcaag   1259
     IleThrThrAsnHisSerGlyLeuTyrAlaCysSerValArgAsnSerAlaThrGlyLys

1260 gaaatctccaaatccatgatagtcaaagtctctggtccctgccatggaaaccagacagag   1319
     GluIleSerLysSerMetIleValLysValSerGlyProCysHisGlyAsnGlnThrGlu

1320 tctcattaatggctgccacaatagagacactgagaaaaagaacaggttgataccttcatg   1379
     SerHisEnd

1380 aaattcaagacaaagaagaaaaaggctcaatgttattggactaaataatcaaaaggataa   1439
1440 tgttttcataatttttattggaaaatgtgctgattcttggaatgttttattctccagatt   1499
1500 tatgaacttttttcttcagcaattggtaaagtatacttttgtaaacaaaaattgaaaca   1559
1560 tttgcttttgctctctatctgagtgcccccc  1591
```

It will be appreciated that the instant specification and claims are set forth by way of illustration and not limitation and that various modifications and changes may be made without departing from the spirit and scope of the present invention.

Claims

1. A nucleic acid comprising a base sequence which codes for a peptide sequence, characterized in that the group nucleic acid is a DNA selected from the following group of five sequences, or is a nucleic acid that is hybridizable with any of such five sequences or that codes for a peptide sequence that is substantially the same as a peptide sequence that is coded for by any of such five sequences:

```
        10                  30                  50
CAGCCGTGCTCGAAGCGTTCCTGGAGCCCAAGCTCTCCTCCACAGGTGAAGACAGGGCCA

        70                  90                  110
GCAGGAGACACCATGGGGCACCTCTCAGCCCCACTTCACAGAGTGCGTGTACCCTGGCAG
           MetGlyHisLeuSerAlaProLeuHisArgValArgValProTrpGln

        130                 150                 170
GGGCTTCTGCTCACAGCCTCACTTCTAACCTTCTGGAACCCGCCCACCACTGCCCAGCTC
GlyLeuLeuLeuThrAlaSerLeuLeuThrPheTrpAsnProProThrThrAlaGlnLeu

        190                 210                 230
ACTACTGAATCCATGCCATTCAATGTTGCAGAGGGGAAGGAGGTTCTTCTCCTTGTCCAC
ThrThrGluSerMetProPheAsnValAlaGluGlyLysGluValLeuLeuLeuValHis
        250                 270                 290
AATCTGCCCCAGCAACTTTTTGGCTACAGCTGGTACAAAGGGGAAAGAGTGGATGGCAAC
AsnLeuProGlnGlnLeuPheGlyTyrSerTrpTyrLysGlyGluArgValAspGlyAsn

        310                 330                 350
CGTCAAATTGTAGGATATGCAATAGGAACTCAACAAGCTACCCCAGGGCCCGCAAACAGC
ArgGlnIleValGlyTyrAlaIleGlyThrGlnGlnAlaThrProGlyProAlaAsnSer

        370                 390                 410
GGTCGAGAGACAATATACCCCAATGCATCCCTGCTGATCCAGAACGTCACCCAGAATGAC
GlyArgGluThrIleTyrProAsnAlaSerLeuLeuIleGlnAsnValThrGlnAsnAsp

        430                 450                 470
ACAGGATTCTACACCCTACAAGTCATAAAGTCAGATCTTGTGAATGAAGAAGCAACTGGA
ThrGlyPheTyrThrLeuGlnValIleLysSerAspLeuValAsnGluGluAlaThrGly
```

```
          490                 510                 530
CAGTTCCATGTATACCCGGAGCTGCCCAAGCCCTCCATCTCCAGCAACAACTCCAACCCT
GlnPheHisValTyrProGluLeuProLysProSerIleSerSerAsnAsnSerAsnPro

          550                 570                 590
GTGGAGGACAAGGATGCTGTGGCCTTCACCTGTGAACCTGAGACTCAGGACACAACCTAC
ValGluAspLysAspAlaValAlaPheThrCysGluProGluThrGlnAspThrThrTyr

          610                 630                 650
CTGTGGTGGATAAACAATCAGAGCCTCCCGGTCAGTCCCAGGCTGCAGCTGTCCAATGGC
LeuTrpTrpIleAsnAsnGlnSerLeuProValSerProArgLeuGlnLeuSerAsnGly

          670                 690                 710
AACAGGACCCTCACTCTACTCAGTGTCACAAGGAATGACACAGGACCCTATGAGTGTGAA
AsnArgThrLeuThrLeuLeuSerValThrArgAsnAspThrGlyProTyrGluCysGlu

          730                 750                 770
ATACAGAACCCAGTGAGTGCGAACCGCAGTGACCCAGTCACCTTGAATGTCACCTATGGC
IleGlnAsnProValSerAlaAsnArgSerAspProValThrLeuAsnValThrTyrGly

          790                 810                 830
CCGGACACCCCCACCATTTCCCCTTCAGACACCTATTACCGTCCAGGGGCAAACCTCAGC
ProAspThrProThrIleSerProSerAspThrTyrTyrArgProGlyAlaAsnLeuSer

          850                 870                 890
CTCTCCTGCTATGCAGCCTCTAACCCACCTGCACAGTACTCCTGGCTTATCAATGGAACA
LeuSerCysTyrAlaAlaSerAsnProProAlaGlnTyrSerTrpLeuIleAsnGlyThr

          910                 930                 950
TTCCAGCAAAGCACACAAGAGCTCTTTATCCCTAACATCACTGTGAATAATAGTGGATCC
PheGlnGlnSerThrGlnGluLeuPheIleProAsnIleThrValAsnAsnSerGlySer

          970                 990                1010
TATACCTGCCACGCCAATAACTCAGTCACTGGCTGCAACAGGACCACAGTCAAGACGATC
TyrThrCysHisAlaAsnAsnSerValThrGlyCysAsnArgThrThrValLysThrIle
```

```
      1030                1050                1070
ATAGTCACTGATAATGCTCTACCACAAGAAAATGGCCTCTCACCTGGGGCCATTGCTGGC
IleValThrAspAsnAlaLeuProGlnGluAsnGlyLeuSerProGlyAlaIleAlaGly

      1090                1110                1130
ATTGTGATTGGAGTAGTGGCCCTGGTTGCTCTGATAGCAGTAGCCCTGGCATGTTTTCTG
IleValIleGlyValValAlaLeuValAlaLeuIleAlaValAlaLeuAlaCysPheLeu

      1150                1170                1190
CATTTCGGGAAGACCGGCAGGGCAAGCGACCAGCGTGATCTCACAGAGCACAAACCCTCA
HisPheGlyLysThrGlyArgAlaSerAspGlnArgAspLeuThrGluHisLysProSer

      1210                1230                1250
GTCTCCAACCACACTCAGGACCACTCCAATGACCCACCTAACAAGATGAATGAAGTTACT
ValSerAsnHisThrGlnAspHisSerAsnAspProProAsnLysMetAsnGluValThr

      1270                1290                1310
TATTCTACCCTGAACTTTGAAGCCCAGCAACCCACACAACCAACTTCAGCCTCCCCATCC
TyrSerThrLeuAsnPheGluAlaGlnGlnProThrGlnProThrSerAlaSerProSer

      1330                1350                1370
CTAACAGCCACAGAAATAATTTATTCAGAAGTAAAAAAAGCAGTAATGAAACCTGTCCTGC
LeuThrAlaThrGluIleIleTyrSerGluValLysLysGln

      1390                1410                1430
TCACTGCAGTGCTGATGTATTTCAAGTCTCTCACCCTCATCACTAGGAGATTCCTTTCCC

      1450                1470                1490
CTGTAGGGTAGAGGGGTGGGGACAGAAACAACTTTCTCCTACTCTTCCTTCCTAATAGGC

      1510                1530                1550
ATCTCCAGGCTGCCTGGTCACTGCCCCTCTCTCAGTGTCAATAGATGAAAGTACATTGGG

      1570                1590                1610
AGTCTGTAGGAAACCCAACCTTCTTGTCATTGAAATTTGGCAAAGCTGACTTTGGGAAAG
```

1630 1650 1670

AGGGACCAGAACTTCCCCTCCCTTCCCCTTTTCCCAACCTGGACTTGTTTTAAACTTGCC

1690 1710 1730

TGTTCAGAGCACTCATTCCTTCCCACCCCCAGTCCTGTCCTATCACTCTAATTCGGATTT

1750 1770 1790

GCCATAGCCTTGAGGTTATGTCCTTTTCCATTAAGTACATGTGCCAGGAAACAGCGAGAG

1810 1830 1850

AGAGAAAGTAAACGGCAGTAATGCTTCTCCTATTTCTCCAAAGCCTTGTGTGAACTAGCA

1870 1890 1910

AAGAGAAGAAAATCAAATATATAACCAATAGTGAAATGCCACAGGTTTGTCCACTGTCAG

1930 1950 1970

GGTTGTCTACCTGTAGGATCAGGGTCTAAGCACCTTGGTGCTTAGCTAGAATACCACCTA

1990 2010 2030

ATCCTTCTGGCAAGCCTGTCTTCAGAGAACCCACTAGAAGCAACTAGGAAAAATCACTTG

2050 2070 2090

CCAAAATCCAAGGCAATTCCTGATGGAAAATGCAAAAGCACATATATGTTTTAATATCTT

2110 2130 2150

TATGGGCTCTGTTCAAGGCAGTGCTGAGAGGGAGGGGTTATAGCTTCAGGAGGGAACCAG

2170 2190 2210

CTTCTGATAAACACAATCTGCTAGGAACTTGGGAAAGGAATCAGAGAGCTGCCCTTCAGC

2230 2250 2270

GATTATTTAAATTGTTAAAGAATACACAATTTGGGGTATTGGGATTTTTCTCCTTTTCTC

2290 2310 2330

TGAGACATTCCACCATTTTAATTTTTGTAACTGCTTATTTATGTGAAAAGGGTTATTTTT

2350 2370 2390

ACTTAGCTTAGCTATGTCAGCCAATCCGATTGCCTTAGGTGAAAGAAACCACCGAAATCC

2410 2430 2450

CTCAGGTCCCTTGGTCAGGAGCCTCTCAAGATTTTTTTTGTCAGAGGCTCCAAATAGAAA

2470 2490 2510

ATAAGAAAAGGTTTTCTTCATTCATGGCTAGAGCTAGATTTAACTCAGTTTCTAGGCACC

2530 2550 2570

TCAGACCAATCATCAACTACCATTCTATTCCATGTTTGCACCTGTGCATTTTCTGTTTGC

2590 2610 2630

CCCCATTCACTTTGTCAGGAAACCTTGGCCTCTGCTAAGGTGTATTTGGTCCTTGAGAAG

2650 2670 2690

TGGGAGCACCCTACAGGGACACTATCACTCATGCTGGTGGCATTGTTTACAGCTAGAAAG

2710 2730 2750

CTGCACTGGTGCTAATGCCCCTTGGGAAATGGGGCTGTGAGGAGGAGGATTATAACTTAG

2770 2790 2810

GCCTAGCCTCTTTTAACAGCCTCTGAAATTTATCTTTTCTTCTATGGGGTCTATAAATGT

2830 2850 2870

ATCTTATAATAAAAAGGAAGGACAGGAGGAAGACAGGCAAATGTACTTCTCACCCAGTCT

2890 2910 2930

TCTACACAGATGGAATCTCTTTGGGGCTAAGAGAAAGGTTTTATTCTATATTGCTTACCT

2950 2970 2990

GATCTCATGTTAGGCCTAAGAGGCTTTCTCCAGGAGGATTAGCTTGGAGTTCTCTATACT

3010 3030 3050

CAGGTACCTCTTTCAGGGTTTTCTAACCCTGACACGGACTGTGCATACTTTCCCTCATCC

3070 3090 3110

ATGCTGTGCTGTGTTATTTAATTTTTCCTGGCTAAGATCATGTCTGAATTATGTATGAAA

3130 3150 3170

ATTATTCTATGTTTTTATAATAAAAATAATATATCAGACATCGAAAAAAAAAA,

(2)

```
          10                  30                  50
          .                   .                   .
CAGCCGTGCTCGAAGCGTTCCTGGAGCCCAAGCTCTCCTCCACAGGTGAAGACAGGGCCA

          70                  90                  110
          .                   .                   .
GCAGGAGACACCATGGGGCACCTCTCAGCCCCACTTCACAGAGTGCGTGTACCCTGGCAG
             MetGlyHisLeuSerAlaProLeuHisArgValArgValProTrpGln

        130                 150                 170
         .                   .                   .
GGGCTTCTGCTCACAGCCTCACTTCTAACCTTCTGGAACCCGCCCACCACTGCCCAGCTC
GlyLeuLeuLeuThrAlaSerLeuLeuThrPheTrpAsnProProThrThrAlaGlnLeu

        190                 210                 230
         .                   .                   .
ACTACTGAATCCATGCCATTCAATGTTGCAGAGGGGAAGGAGGTTCTTCTCCTTGTCCAC
ThrThrGluSerMetProPheAsnValAlaGluGlyLysGluValLeuLeuLeuValHis

        250                 270                 290
         .                   .                   .
AATCTGCCCCAGCAACTTTTTGGCTACAGCTGGTACAAAGGGGAAAGAGTGGATGGCAAC
AsnLeuProGlnGlnLeuPheGlyTyrSerTrpTyrLysGlyGluArgValAspGlyAsn

        310                 330                 350
         .                   .                   .
CGTCAAATTGTAGGATATGCAATAGGAACTCAACAAGCTACCCCAGGGCCCGCAAACAGC
ArgGlnIleValGlyTyrAlaIleGlyThrGlnGlnAlaThrProGlyProAlaAsnSer

        370                 390                 410
         .                   .                   .
GGTCGAGAGACAATATACCCCAATGCATCCCTGCTGATCCAGAACGTCACCCAGAATGAC
GlyArgGluThrIleTyrProAsnAlaSerLeuLeuIleGlnAsnValThrGlnAsnAsp
```

```
          430                 450                 470
ACAGGATTCTACACCCTACAAGTCATAAAGTCAGATCTTGTGAATGAAGAAGCAACTGGA
ThrGlyPheTyrThrLeuGlnValIleLysSerAspLeuValAsnGluGluAlaThrGly

          490                 510                 530
CAGTTCCATGTATACCCGGAGCTGCCCAAGCCCTCCATCTCCAGCAACAACTCCAACCCT
GlnPheHisValTyrProGluLeuProLysProSerIleSerSerAsnAsnSerAsnPro

          550                 570                 590
GTGGAGGACAAGGATGCTGTGGCCTTCACCTGTGAACCTGAGACTCAGGACACAACCTAC
ValGluAspLysAspAlaValAlaPheThrCysGluProGluThrGlnAspThrThrTyr

          610                 630                 650
CTGTGGTGGATAAACAATCAGAGCCTCCCGGTCAGTCCCAGGCTGCAGCTGTCCAATGGC
LeuTrpTrpIleAsnAsnGlnSerLeuProValSerProArgLeuGlnLeuSerAsnGly

          670                 690                 710
AACAGGACCCTCACTCTACTCAGTGTCACAAGGAATGACACAGGACCCTATGAGTGTGAA
AsnArgThrLeuThrLeuLeuSerValThrArgAsnAspThrGlyProTyrGluCysGlu

          730                 750                 770
ATACAGAACCCAGTGAGTGCGAACCGCAGTGACCCAGTCACCTTGAATGTCACCTATGGC
IleGlnAsnProValSerAlaAsnArgSerAspProValThrLeuAsnValThrTyrGly

          790                 810                 830
CCGGACACCCCCACCATTTCCCCTTCAGACACCTATTACCGTCCAGGGGCAAACCTCAGC
ProAspThrProThrIleSerProSerAspThrTyrTyrArgProGlyAlaAsnLeuSer
```

```
          850                 870                 890
CTCTCCTGCTATGCAGCCTCTAACCCACCTGCACAGTACTCCTGGCTTATCAATGGAACA
LeuSerCysTyrAlaAlaSerAsnProProAlaGlnTyrSerTrpLeuIleAsnGlyThr

          910                 930                 950
TTCCAGCAAAGCACACAAGAGCTCTTTATCCCTAACATCACTGTGAATAATAGTGGATCC
PheGlnGlnSerThrGlnGluLeuPheIleProAsnIleThrValAsnAsnSerGlySer

          970                 990                1010
TATACCTGCCACGCCAATAACTCAGTCACTGGCTGCAACAGGACCACAGTCAAGACGATC
TyrThrCysHisAlaAsnAsnSerValThrGlyCysAsnArgThrThrValLysThrIle

         1030                1050                1070
ATAGTCACTGAGCTAAGTCCAGTAGTAGCAAAGCCCCAAATCAAAGCCAGCAAGACCACA
IleValThrGluLeuSerProValValAlaLysProGlnIleLysAlaSerLysThrThr

         1090                1110                1130
GTCACAGGAGATAAGGACTCTGTGAACCTGACCTGCTCCACAAATGACACTGGAATCTCC
ValThrGlyAspLysAspSerValAsnLeuThrCysSerThrAsnAspThrGlyIleSer

         1150                1170                1190
ATCCGTTGGTTCTTCAAAAACCAGAGTCTCCCGTCCTCGGAGAGGATGAAGCTGTCCCAG
IleArgTrpPhePheLysAsnGlnSerLeuProSerSerGluArgMetLysLeuSerGln

         1210                1230                1250
GGCAACACCACCCTCAGCATAAACCCTGTCAAGAGGGAGGATGCTGGGACGTATTGGTGT
GlyAsnThrThrLeuSerIleAsnProValLysArgGluAspAlaGlyThrTyrTrpCys
```

```
      1270                1290                1310
GAGGTCTTCAACCCAATCAGTAAGAACCAAAGCGACCCCATCATGCTGAACGTAAACTAT
GluValPheAsnProIleSerLysAsnGlnSerAspProIleMetLeuAsnValAsnTyr

      1330                1350                1370
AATGCTCTACCACAAGAAAATGGCCTCTCACCTGGGGCCATTGCTGGCATTGTGATTGGA
AsnAlaLeuProGlnGluAsnGlyLeuSerProGlyAlaIleAlaGlyIleValIleGly

      1390                1410                1430
GTAGTGGCCCTGGTTGCTCTGATAGCAGTAGCCCTGGCATGTTTTCTGCATTTCGGGAAG
ValValAlaLeuValAlaLeuIleAlaValAlaLeuAlaCysPheLeuHisPheGlyLys

      1450                1470                1490
ACCGGCAGCTCAGGACCACTCCAATGACCCACCTAACAAGATGAATGAAGTTACTTATTC
ThrGlySerSerGlyProLeuGln

      1510                1530                1550
TACCCTGAACTTTGAAGCCCAGCAACCCACACAACCAACTTCAGCCTCCCCATCCCTAAC

      1570                1590                1610
AGCCACAGAAATAATTTATTCAGAAGTAAAAAAAGCAGTAATGAAACCTGAAAAAAAAAAA

      1630
AAAAAAAAAA
```

(3)

```
          10                  30                  50
CAGCCGTGCTCGAAGCGTTCCTGGAGCCCAAGCTCTCCTCCACAGGTGAAGACAGGGCCA

          70                  90                  110
GCAGGAGACACCATGGGGCACCTCTCAGCCCCACTTCACAGAGTGCGTGTACCCTGGCAG
              MetGlyHisLeuSerAlaProLeuHisArgValArgValProTrpGln

         130                 150                 170
GGGCTTCTGCTCACAGCCTCACTTCTAACCTTCTGGAACCCGCCCACCACTGCCCAGCTC
GlyLeuLeuLeuThrAlaSerLeuLeuThrPheTrpAsnProProThrThrAlaGlnLeu

         190                 210                 230
ACTACTGAATCCATGCCATTCAATGTTGCAGAGGGGAAGGAGGTTCTTCTCCTTGTCCAC
ThrThrGluSerMetProPheAsnValAlaGluGlyLysGluValLeuLeuLeuValHis

         250                 270                 290
AATCTGCCCCAGCAACTTTTTGGCTACAGCTGGTACAAAGGGGAAAGAGTGGATGGCAAC
AsnLeuProGlnGlnLeuPheGlyTyrSerTrpTyrLysGlyGluArgValAspGlyAsn

         310                 330                 350
CGTCAAATTGTAGGATATGCAATAGGAACTCAACAAGCTACCCCAGGGCCCGCAAACAGC
ArgGlnIleValGlyTyrAlaIleGlyThrGlnGlnAlaThrProGlyProAlaAsnSer

         370                 390                 410
GGTCGAGAGACAATATACCCCAATGCATCCCTGCTGATCCAGAACGTCACCCAGAATGAC
GlyArgGluThrIleTyrProAsnAlaSerLeuLeuIleGlnAsnValThrGlnAsnAsp

         430                 450                 470
ACAGGATTCTACACCCTACAAGTCATAAAGTCAGATCTTGTGAATGAAGAAGCAACTGGA
ThrGlyPheTyrThrLeuGlnValIleLysSerAspLeuValAsnGluGluAlaThrGly
```

```
          490                 510                 530
CAGTTCCATGTATACCCGGAGCTGCCCAAGCCCTCCATCTCCAGCAACAACTCCAACCCT
GlnPheHisValTyrProGluLeuProLysProSerIleSerSerAsnAsnSerAsnPro

          550                 570                 590
GTGGAGGACAAGGATGCTGTGGCCTTCACCTGTGAACCTGAGACTCAGGACACAACCTAC
ValGluAspLysAspAlaValAlaPheThrCysGluProGluThrGlnAspThrThrTyr

          610                 630                 650
CTGTGGTGGATAAACAATCAGAGCCTCCCGGTCAGTCCCAGGCTGCAGCTGTCCAATGGC
LeuTrpTrpIleAsnAsnGlnSerLeuProValSerProArgLeuGlnLeuSerAsnGly

          670                 690                 710
AACAGGACCCTCACTCTACTCAGTGTCACAAGGAATGACACAGGACCCTATGAGTGTGAA
AsnArgThrLeuThrLeuLeuSerValThrArgAsnAspThrGlyProTyrGluCysGlu

          730                 750                 770
ATACAGAACCCAGTGAGTGCGAACCGCAGTGACCCAGTCACCTTGAATGTCACCTATGGC
IleGlnAsnProValSerAlaAsnArgSerAspProValThrLeuAsnValThrTyrGly

          790                 810                 830
CCGGACACCCCCACCATTTCCCCTTCAGACACCTATTACCGTCCAGGGGCAAACCTCAGC
ProAspThrProThrIleSerProSerAspThrTyrTyrArgProGlyAlaAsnLeuSer

          850                 870                 890
CTCTCCTGCTATGCAGCCTCTAACCCACCTGCACAGTACTCCTGGCTTATCAATGGAACA
LeuSerCysTyrAlaAlaSerAsnProProAlaGlnTyrSerTrpLeuIleAsnGlyThr

          910                 930                 950
TTCCAGCAAAGCACACAAGAGCTCTTTATCCCTAACATCACTGTGAATAATAGTGGATCC
PheGlnGlnSerThrGlnGluLeuPheIleProAsnIleThrValAsnAsnSerGlySer

          970                 990                 1010
TATACCTGCCACGCCAATAACTCAGTCACTGGCTGCAACAGGACCACAGTCAAGACGATC
TyrThrCysHisAlaAsnAsnSerValThrGlyCysAsnArgThrThrValLysThrIle
```

```
      1030                1050                1070
ATAGTCACTGATAATGCTCTACCACAAGAAAATGGCCTCTCACCTGGGGCCATTGCTGGC
IleValThrAspAsnAlaLeuProGlnGluAsnGlyLeuSerProGlyAlaIleAlaGly

      1090                1110                1130
ATTGTGATTGGAGTAGTGGCCCTGGTTGCTCTGATAGCAGTAGCCCTGGCATGTTTTCTG
IleValIleGlyValValAlaLeuValAlaLeuIleAlaValAlaLeuAlaCysPheLeu

      1150                1170                1190
CATTTCGGGAAGACCGGCAGCTCAGGACCACTCCAATGACCCACCTAACAAGATGAATGA
HisPheGlyLysThrGlySerSerGlyProLeuGln

      1210                1230                1250
AGTTACTTATTCTACCCTGAACTTTGAAGCCCAGCAACCCACACAACCAACTTCAGCCTC

      1270                1290                1310
CCCATCCCTAACAGCCACAGAAATAATTTATTCAGAAGTAAAAAAAGCAGTAATGAAACCT

      1330
GAAAAAAAAAAAAAAAAAA ;
```

(4)

```
   1  acagcacagctgacagccgtactcaggaagcttctggatcctaggcttatctccacagag    60

  61  gagaacacacaagcagcagagaccatggggccccctctcagcccctccctgcacacacctc   120
                                    MetGlyProLeuSerAlaProProCysThrHisLeu

 121  atcacttggaaggggtcctgctcacagcatcacttttaaacttctggaatccgcccaca     180
      IleThrTrpLysGlyValLeuLeuThrAlaSerLeuLeuAsnPheTrpAsnProProThr

 181  actgcccaagtcacgattgaagcccagccacccaaagtttctgaggggaaggatgttctt    240
      ThrAlaGlnValThrIleGluAlaGlnProProLysValSerGluGlyLysAspValLeu

 241  ctacttgtccacaatttgccccagaatcttgctggctacatttggtacaaagggcaaatg    300
      LeuLeuValHisAsnLeuProGlnAsnLeuAlaGlyTyrIleTrpTyrLysGlyGlnMet

 301  acatacgtctaccattacattacatcatatgtagtagacggtcaaagaattatatatggg    360
      ThrTyrValTyrHisTyrIleThrSerTyrValValAspGlyGlnArgIleIleTyrGly

 361  cctgcatacagtggaagagaaagagtatattccaatgcatccctgctgatccagaatgtc    420
      ProAlaTyrSerGlyArgGluArgValTyrSerAsnAlaSerLeuLeuIleGlnAsnVal

 421  acgcaggaggatgcaggatcctacaccttacacatcataaagcgacgcgatgggactgga    480
      ThrGlnGluAspAlaGlySerTyrThrLeuHisIleIleLysArgArgAspGlyThrGly

 481  ggagtaactggacatttcaccttcaccttacacctggagactcccaagccctccatctcc    540
      GlyValThrGlyHisPheThrPheThrLeuHisLeuGluThrProLysProSerIleSer

 541  agcagcaacttaaatcccagggaggccatggaggctgtgatcttaacctgtgatcctgcg    600
      SerSerAsnLeuAsnProArgGluAlaMetGluAlaValIleLeuThrCysAspProAla

 601  actccagccgcaagctaccagtggtggatgaatggtcagagcctccctatgactcacagg    660
      ThrProAlaAlaSerTyrGlnTrpTrpMetAsnGlyGlnSerLeuProMetThrHisArg

 661  ttgcagctgtccaaaaccaacaggaccctctttatatttggtgtcacaaagtatattgca    720
      LeuGlnLeuSerLysThrAsnArgThrLeuPheIlePheGlyValThrLysTyrIleAla

 721  ggaccctatgaatgtgaaatacggaacccagtgagtgccagccgcagtgacccagtcacc    780
      GlyProTyrGluCysGluIleArgAsnProValSerAlaSerArgSerAspProValThr

 781  ctgaatctcctcccaaagctgtccaagccctacatcacaatcaacaacttaaaccccaga    840
      LeuAsnLeuLeuProLysLeuSerLysProTyrIleThrIleAsnAsnLeuAsnProArg

 841  gagaataaggatgtcttaaccttcacctgtgaacctaagagtgagaactacacctacatt    900
      GluAsnLysAspValLeuThrPheThrCysGluProLysSerGluAsnTyrThrTyrIle

 901  tggtggctaaatggtcagagcctccctgtcagtcccagggtaaagcgacccattgaaaac    960
      TrpTrpLeuAsnGlyGlnSerLeuProValSerProArgValLysArgProIleGluAsn

 961  aggatcctcattctacccaatgtcacgagaaatgaaacaggaccttatcaatgtgaaata   1020
      ArgIleLeuIleLeuProAsnValThrArgAsnGluThrGlyProTyrGlnCysGluIle

1021  cgggaccgatatggtggcatccgcagtgacccagtcaccctgaatgtcctctatggtcca   1080
      ArgAspArgTyrGlyGlyIleArgSerAspProValThrLeuAsnValLeuTyrGlyPro
```

```
1081  gacctccccagcatttacccttcattcacctattaccgttcaggagaaaacctctacttt  1140
      AspLeuProSerIleTyrProSerPheThrTyrTyrArgSerGlyGluAsnLeuTyrPhe

1141  tcctgcttcggtgagtctaacccacgggcacaatattcttggacaattaatgggaagttt  1200
      SerCysPheGlyGluSerAsnProArgAlaGlnTyrSerTrpThrIleAsnGlyLysPhe

1201  cagctatcaggacaaaagctctctatcccccaaataactacaaagcatagtgggctctat  1260
      GlnLeuSerGlyGlnLysLeuSerIleProGlnIleThrThrLysHisSerGlyLeuTyr

1261  gcttgctctgttcgtaactcagccactggcaaggaaagctccaaatccatcacagtcaaa  1320
      AlaCysSerValArgAsnSerAlaThrGlyLysGluSerSerLysSerIleThrValLys

1321  gtctctgactggatattaccctgaattctactagttcctccaattccattttctcccatg  1380
      ValSerAspTrpIleLeuProEnd

1381  gaatcacgaagagcaagacccactctgttccagaagccctataatctggaggtggacaac  1440
1441  tcgatgtaaatttcatgggaaaaccccttgtacctgacatgtgagccactcagaactcacc  1500
1501  aaaatgttcgacaccataacaacagctactcaaactgtaaaccaggataagaagttgatg  1560
1561  acttcacactgtggacagtttttcaaagatgtcataacaagactccccatcatgacaagg  1620
1621  ctccaccctctactgtctgctcatgcctgcctctttcacttggcaggataatgcagtcat  1680
1681  tagaatttcacatgtagtagcttctgagggtaacaacagagtgtcagatatgtcatctca  1740
1741  acctcaaacttttacgtaacatctcagggaaatgtggctctctccatcttgcatacaggg  1800
1801  ctcccaatagaaatgaacacagagatattgcctgtgtgtttgcagagaagatggtttcta  1860
1861  taaagagtaggaaagctgaaattatagtagagtctcctttaaatgcacattgtgtggatg  1920
1921  gctctcaccatttcctaagagatacagtgtaaaaacgtgacagtaatactgattctagca  1980
1981  gaataaacatgtaccacatttgcaaaaaa                                  2010
```

and

(5)

```
    1  gggtggatcctaggctcatctccataggggagaacacacatacagcagagaccatggga   59
                                                              MetGly

   60  cccctctcagcccctccctgcactcagcacatcacctggaaggggctcctgctcacagca  119
       ProLeuSerAlaProProCysThrGlnHisIleThrTrpLysGlyLeuLeuLeuThrAla

  120  tcacttttaaacttctggaacctgcccaccactgcccaagtaataattgaagcccagcca  179
       SerLeuLeuAsnPheTrpAsnLeuProThrThrAlaGlnValIleIleGluAlaGlnPro

  180  cccaaagtttctgaggggaaggatgttcttctacttgtccacaatttgccccagaatctt  239
       ProLysValSerGluGlyLysAspValLeuLeuLeuValHisAsnLeuProGlnAsnLeu

  240  actggctacatctggtacaaagggcaaatgacggacctctaccattacattacatcatat  299
       ThrGlyTyrIleTrpTyrLysGlyGlnMetThrAspLeuTyrHisTyrIleThrSerTyr

  300  gtagtagacggtcaaattatatatgggcctgcctacagtggacgagaaacagtatattcc  359
       ValValAspGlyGlnIleIleTyrGlyProAlaTyrSerGlyArgGluThrValTyrSer

  360  aatgcatccctgctgatccagaatgtcacacaggaggatgcaggatcctacaccttacac  419
       AsnAlaSerLeuLeuIleGlnAsnValThrGlnGluAspAlaGlySerTyrThrLeuHis

  420  atcataaagcgaggcgatgggactggaggagtaactggatatttcactgtcaccttatac  479
       IleIleLysArgGlyAspGlyThrGlyGlyValThrGlyTyrPheThrValThrLeuTyr

  480  tcggagactcccaagcgctccatctccagcagcaacttaaaccccagggaggtcatggag  539
       SerGluThrProLysArgSerIleSerSerSerAsnLeuAsnProArgGluValMetGlu

  540  gctgtgcgcttaatctgtgatcctgagactccggatgcaagctacctgtggttgctgaat  599
       AlaValArgLeuIleCysAspProGluThrProAspAlaSerTyrLeuTrpLeuLeuAsn

  600  ggtcagaacctccctatgactcacaggttgcagctgtccaaaaccaacaggaccctctat  659
       GlyGlnAsnLeuProMetThrHisArgLeuGlnLeuSerLysThrAsnArgThrLeuTyr

  660  ctatttggtgtcacaaagtatattgcagggccctatgaatgtgaaatacggaggggagtg  719
       LeuPheGlyValThrLysTyrIleAlaGlyProTyrGluCysGluIleArgArgGlyVal

  720  agtgccagccgcagtgacccagtcaccctgaatctcctcccgaagctgcccatgccttac  779
       SerAlaSerArgSerAspProValThrLeuAsnLeuLeuProLysLeuProMetProTyr

  780  atcaccatcaacaacttaaaccccagggagaagaaggatgtgttagccttcacctgtgaa  839
       IleThrIleAsnAsnLeuAsnProArgGluLysLysAspValLeuAlaPheThrCysGlu

  840  cctaagagtcggaactacacctacatttggtggctaaatggtcagagcctcccggtcagt  899
       ProLysSerArgAsnTyrThrTyrIleTrpTrpLeuAsnGlyGlnSerLeuProValSer

  900  ccgagggtaaagcgacccattgaaaacaggatactcattctacccagtgtcacgagaaat  959
       ProArgValLysArgProIleGluAsnArgIleLeuIleLeuProSerValThrArgAsn

  960  gaaacaggaccctatcaatgtgaaatacgggaccgatatggtggcatccgcagtaaccca 1019
       GluThrGlyProTyrGlnCysGluIleArgAspArgTyrGlyGlyIleArgSerAsnPro
```

```
1020  gtcaccctgaatgtcctctatggtccagacctccccagaatttacccttacttcacctat  1079
      ValThrLeuAsnValLeuTyrGlyProAspLeuProArgIleTyrProTyrPheThrTyr

1080  taccgttcaggagaaaacctcgacttgtcctgctttgcggactctaacccaccggcagag  1139
      TyrArgSerGlyGluAsnLeuAspLeuSerCysPheAlaAspSerAsnProProAlaGlu

1140  tatttttggacaattaatgggaagtttcagctatcaggacaaaagctctttatcccccaa  1199
      TyrPheTrpThrIleAsnGlyLysPheGlnLeuSerGlyGlnLysLeuPheIleProGln

1200  attactacaaatcatagcgggctctatgcttgctctgttcgtaactcagccactggcaag  1259
      IleThrThrAsnHisSerGlyLeuTyrAlaCysSerValArgAsnSerAlaThrGlyLys

1260  gaaatctccaaatccatgatagtcaaagtctctggtccctgccatggaaaccagacagag  1319
      GluIleSerLysSerMetIleValLysValSerGlyProCysHisGlyAsnGlnThrGlu

1320  tctcattaatggctgccacaatagagacactgagaaaaagaacaggttgataccttcatg  1379
      SerHisEnd

1380  aaattcaagacaaagaagaaaaaggctcaatgttattggactaaataatcaaaaggataa  1439
1440  tgttttcataatttttattggaaaatgtgctgattcttggaatgttttattctccagatt  1499
1500  tatgaacttttttcttcagcaattggtaaagtatacttttgtaaacaaaaattgaaaca  1559
1560  tttgcttttgctctctatctgagtgcccccc  1591
```

2. A nucleic acid comprising a base sequence which codes for the protein CEA-(e), characterized in that it is DNA sequence (1) of claim 1.

3. A nucleic acid comprising a base sequence which codes for the protein CEA-(f), characterized in that it is DNA sequence (2) of claim 1.

4. A nucleic acid comprising a base sequence which codes for the protein CEA-(g), characterized in that it is DNA sequence (3) of claim 1.

5. A nucleic acid comprising a base sequence which codes for the protein KGCEA1, characterized in that it is DNA sequence (4) of claim 1.

6. A nucleic acid comprising a base sequence which codes for the protein KGCEA2, characterized in that it is DNA sequence (5) of claim 1.

7. A replicable recombinant cloning vehicle having an insert comprising a nucleic acid of any one of claims 1-6.

8. A cell that is transfected, infected or injected with a recombinant cloning vehicle of claim 7.

9. A protein characterized by having an amino acid sequence coded by a nucleic acid of any one of claims 1-6, or a polypeptide or peptide fragment thereof having no less than five amino acids.

10. An antibody prepared against a protein, polypeptide, or peptide of claim 9.


TM1(CEA(c))
10
12
14
16
18
I
IIa
TM2(CEA(e))
10
12
16
18
I
20
TM3(CEA(f))
10
12
14
16
24
26
I
IIa
22
TM4(CEA(g))
10
12
16
24
26
I
22
20

FIG.1

(19) Europäisches Patentamt
European Patent Office
Office européen des brevets

(11) Publication number: **0 346 710 A3**

(12) # EUROPEAN PATENT APPLICATION

(21) Application number: **89110096.8**

(22) Date of filing: **03.06.89**

(51) Int. Cl.[5]: **C12N 15/00**, C12N 5/00, C07K 13/00, A61K 39/395

(30) Priority: **16.06.88 US 207678**
21.11.88 US 274107

(43) Date of publication of application:
20.12.89 Bulletin 89/51

(84) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI NL SE

(88) Date of deferred publication of the search report:
11.09.91 Bulletin 91/37

(71) Applicant: **MOLECULAR DIAGNOSTICS, INC.**
400 Morgan Lane
West Haven, CT 06516(US)

(72) Inventor: **Barnett, Thomas R., Dr.**
27 Jeffrey Road
East Haven, CT 06513(US)
Inventor: **Elting, James J., Dr.**
5 Heatherwood Drive
Madison, CT 06443(US)
Inventor: **Kamarck, Michael E.**
86 Russell Road
Bethany, CT 06525(US)
Inventor: **Kretschmer, Axel, Dr.**
Richard-Zörner-Strasse 32
W-5060 Bergisch Gladbach 1(DE)

(74) Representative: **Adrian, Albert, Dr. et al**
c/o BAYER AG Konzernverwaltung RP
Patentabteilung
W-5090 Leverkusen 1, Bayerwerk(DE)

(54) **cDNAs coding for members of the carcinoembryonic antigen family.**

(57) A nucleic acid comprising a base sequence which codes for a CEA family member peptide sequence or nucleic acids having a base sequence hybridizable therewith, replicable recombinant cloning vehicles having an insert comprising such nucleic acid, cells transfected, infected or injected with such cloning vehicles, polypeptides expressed by such cells, synthetic peptides derived from the coding sequence of CEA family member nucleic acids, antibody preparations specific for such polypeptides, immunoassays for detecting CEA family members using such antibody preparations and nucleic acid hybridization methods for detecting CEA family member nucleic acid sequences using a nucleic acid probe comprising the above described nucleic acid.

EP 0 346 710 A3

European Patent Office

EUROPEAN SEARCH REPORT

Application Number

EP 89 11 0096

DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
X	BIOCHEM. BIOPHYS. RES. COMMUN., vol. 142, no. 2, 30th January 1987, pages 511-518; R. OIKAWA et al.: "Primary structure of human carcinoembryonic antigen (CEA) deduced from cDNA sequence" * The whole document *	1,6-10	C 12 N 15/00 C 12 N 5/00 C 07 K 13/00 A 61 K 39/395
X	MOL. CELL. BIOL., vol. 7, 1987, page 3221-3230; R. BEAUCHEMIN et al.: "Isolation and characterization of full-lenght functional cDNA clones for human carcinoembryonic antigen" * The whole document *	1,6-10	
P,X	PROC. NATL. ACAD. SCI. USA, vol. 85, September 1988, pages 6959-6963; Y. HINODA et al.: "Molecular cloning of a cDNA coding biliary glycoprotein I: primary structure of a glycoprotein immunologically crossreactive with carcinoembryonic antigen" * The whole document *	1,6-10	
P,X	GENE, vol. 71, no. 2, November 1988, pages 439-449; B.C. ROONEY et al.: "Molecular cloning of a cDNA for human pregnancy-specific B1-glycoprotein: homology with human carcinoembryonic antigen and related proteins" * The whole document *	1	
A	EP-A-0 212 880 (STATE OF ISRAEL)		

TECHNICAL FIELDS SEARCHED (Int. Cl.5)

C 07 K
C 12 N

The present search report has been drawn up for all claims

Place of search	Date of completion of search	Examiner
The Hague	17 May 91	NAUCHE S.A.

CATEGORY OF CITED DOCUMENTS

X: particularly relevant if taken alone
Y: particularly relevant if combined with another document of the same catagory
A: technological background
O: non-written disclosure
P: intermediate document
T: theory or principle underlying the invention

E: earlier patent document, but published on, or after the filing date
D: document cited in the application
L: document cited for other reasons

&: member of the same patent family, corresponding document